

PUBLICATIONS IN ENGLISH (Akira Nakagawara, M.D., Ph.D.)

Original Publications

1. **Nakagawara, A.**, Takeshige, K., Minakami, S. Induction of a phagocytosis-like metabolic pattern in polymorphonuclear leucocytes by cytochalasin E. *Exp. Cell Res.* 87:392-394, 1974.
2. Takeshige, S., **Nakagawara, A.**, Minakami, S. Metabolic pattern of polymorphonuclear leucocytes induced by trypsin-digested microsomes. *J. Biochem.* 77:559-566, 1975.
3. **Nakagawara, A.**, Takeshige, K., Minakami, S. Effect of cyclic nucleotides on the cyanide-insensitive respiration of polymorphonuclear leucocytes. *J. Biochem.* 77:567-573, 1975.
4. **Nakagawara, A.**, Minakami, S. Generation of superoxide anions by leucocytes treated with cytochalasin E. *Biochem. Biophys. Res. Commun.* 64:760-767, 1975.
5. **Nakagawara, A.**, Kakinuma, K., Shin, H., Miyazaki, S., Minakami, S. Lack of cytochalasin E induced superoxide release by polymorphonuclear leucocytes of patients with chronic granulomatous disease. A diagnostic test. *Clin. Chim. Acta* 70:133-137, 1976.
6. **Nakagawara, A.**, Shibata, Y., Takeshige, K., Minakami, S. Action of cytochalasin E on polymorphonuclear leucocytes of guinea pig peritoneal exudates. *Exp. Cell Res.* 101:225-234, 1976.
7. Nakamura, M., **Nakagawara, A.**, Minakami, S. Mimicking of phagocytotic metabolism by leucocytes suspended in deuterium oxide. *Exp. Cell Res.* 102:429-431, 1976.
8. Miyazaki, S., Shin, H., Goya, N., **Nakagawara, A.** Identification of a carrier mother of a female patient with chronic granulomatous disease. *J. Pediatr.* 89:784-786, 1976.
9. **Nakagawara, A.**, Nabi, B.Z.F., Minakami, S. An improved procedure for the diagnosis of chronic granulomatous disease, using concanavalin A and cytochalasin E. *Clin. Chim. Acta* 74:173-176, 1977.
10. Minakami, S., **Nakagawara, A.**, Nakamura, M. Generation and release of superoxide anions by leucocytes. In: Hayaishi, O. (ed): *Biochemical and Medical Aspects of Active Oxygen*. University of Tokyo Press, Tokyo, 1977, pp. 229-246.
11. **Nakagawara, A.**, Minakami, S. Role of cytoskeletal elements in cytochalasin E-induced superoxide production by human polymorphonuclear leucocytes. *Biochim. Biophys. Acta* 584:143-148, 1979.
12. Ikeda, K., Suita, S., **Nakagawara, A.**, Takabayashi, K. Preoperative chemotherapy for initially unresectable hepatoblastoma in children. Survival in two cases. *Arch. Surg.* 114:203-207, 1979.
13. **Nakagawara, A.**, Kayashima, K., Tamada, R., Onoue, K., Ikeda, K., Inokuchi, K. A sensitive and rapid method for determination of superoxide generating activity of blood monocytes and its use as a probe for monocyte *function in cancer patients*. *Gann* 70:829-833, 1979.
14. **Nakagawara, A.**, Ikeda, K., Ohgami, H. Mucoepidermoid tumor of the bronchus in an infant. *J. Pediatr. Surg.* 14:608-609, 1979.
15. **Nakagawara, A.** Simple functional tests for polymorphonuclear leukocytes using cytochalasin E and lectins. In: Kokubun, Y., Kobayashi, N. (eds): *Phagocytosis. Its Physiology and Pathology*. University of Tokyo Press, Tokyo, 1979, pp. 45-52.

16. Miyazaki, S., Shin, H., Goya, N., **Nakagawara, A.** Identification of a carrier mother of a female patient with chronic granulomatous disease. In: Kokubun, Y., Kobayashi, N. (eds): *Phagocytosis. Its Physiology and Pathology*. University of Tokyo Press, Tokyo, 1979, pp. 227-231.
17. Kayashima, K., Onoue, K., **Nakagawara, A.**, Minakami, S. Superoxide anion-generating activities of macrophages as studied by using cytochalasin E and lectins as synergistic stimulants for superoxide release. *Microbiol. Immunol.* 24:449-461, 1980.
18. Goto, S., Ikeda, K., Kume, K., **Nakagawara, A.** Postoperative acetylcholinesterase activity in Hirschsprung's disease. *Z. Kinderchir.* 30:217-224, 1980.
19. **Nakagawara, A.**, Nathan, C.F., Cohn, Z.A. Hydrogen peroxide metabolism in human monocytes during differentiation in vitro. *J. Clin. Invest.* 68:1243-1252, 1981.
20. Ikeda, K., Kimura, N., Suita, S., Hayashida, Y., **Nakagawara, A.**, Kawaoi, A. Pre and postoperative changes of alpha-fetoprotein and human chorionic gonadotropin in hepatoma of children. *Z. Kinderchir.* 34:42-55, 1981.
21. **Nakagawara, A.**, De Santis, N.M., Nogueira, N., Nathan, C.F. Lymphokines enhance the capacity of human monocytes to secrete reactive oxygen intermediates. *J. Clin. Invest.* 70:1042-1048, 1982.
22. **Nakagawara, A.**, Ikeda, K., Hayashida, Y., Tsuneyoshi, M., Enjoji, M., Kawaoi, A. Immunocytochemical identification of human chorionic gonadotropin and alpha-fetoprotein-producing cells of hepatoblastoma associated with precocious puberty. *Virchows Arch [Pathol Anat]* 398:45-51, 1982.
23. Nathan, C.F., **Nakagawara, A.** Role of reactive oxygen intermediates in macrophage killing of intracellular pathogens: A review. In: Mizuno, D., Cohn, Z.A., Takeya, K., Ishida, N. (eds): *Self-defense Mechanisms. Role of Macrophages*. University of Tokyo Press, Tokyo, 1982, pp. 279-294.
24. **Nakagawara, A.**, Nathan, C.F. A simple method for counting adherent cells: Application to cultured human monocytes, macrophages and multinucleated giant cells. *J. Immunol. Meth.* 56:261-268, 1983.
25. Matsuo, S., Ikeda, K., Yakabe, S., **Nakagawara, A.**, Iwashita, A. Histological study of the remnant of porta hepatis in patients with extrahepatic biliary atresia. A computed picture analysis of 30 cases. *Z. Kinderchir.* 39:46-49, 1984.
26. **Nakagawara, A.**, Ikeda, K., Inokuchi, K., Kumashiro, R., Tamada, R. Deficient superoxide-generating activity and its activation of blood monocytes in cancer patients. *Cancer Lett.* 22:157-162, 1984.
27. Yakabe, S., Ikeda, K., Ohgami, H., **Nakagawara, A.**, Matsuo, S., Yae, Y. Clinical significance of lipoprotein-X in congenital biliary atresia. *Z. Kinderchir.* 39:168-170, 1984.
28. **Nakagawara, A.**, Inokuchi, K., Ikeda, K., Kumashiro, R., Tamada, R. Decreased superoxide-generating activity of blood monocytes from patients with hepatic cirrhosis. *Hepato-Gastroenterol.* 31:201-203, 1984.
29. Ishimaru, K., **Nakagawara, A.**, Zaizen, Y., Ikeda, K. Combined treatment with cisplatin and radiation therapy for mouse C1300 neuroblastoma. *Z. Kinderchir.* 40:36-39, 1985.
30. **Nakagawara, A.**, Toyohara, T., Ikeda, K., Nada, O. VMA-negative neuroblastoma in children. *Eur. J. Pediatr.* 144:421, 1985.

31. **Nakagawara, A.**, Ikeda, K., Tsuneyoshi, M., Daimaru, Y., Enjoji, M., Watanabe, I., Iwafuchi, M., Sawada, T. Hepatoblastoma producing both alpha-fetoprotein (AFP) and human chorionic gonadotropin (hCG). Clinicopathological analysis of 4 cases and a review of the literature. *Cancer* 56:144-150, 1985.
32. Matsuo, S., **Nakagawara, A.**, Ikeda, K., Mitsuyama, M., Nomoto, K. Enhanced release of reactive oxygen intermediates by immunologically activated rat Kupffer cells. *Clin. Exp. Immunol.* 59:203-209, 1985.
33. **Nakagawara, A.**, Ikeda, K., Tsuneyoshi, M., Daimaru, Y., Enjoji, M. Malignant pheochromocytoma with ganglioneuroblastomatous elements in a patient with von Recklinghausen's disease. *Cancer* 55:2794-2798, 1985.
34. **Nakagawara, A.**, Toyohara, T., Nada, O., Ikeda, K. Catecholaminergic differentiation associated with S100 protein-positive elements in human neuroblastoma. *Z. Kinderchir.* 41:275-278, 1986.
35. Goto, S., Ikeda, K., **Nakagawara, A.**, Daimaru, Y., Tsuneyoshi, M., Enjoji, M. Renal cell carcinoma in Japanese children. *J. Urol.* 136:1261-1263, 1986
36. **Nakagawara, A.**, Toyohara, T., Ikeda, K., Nada, O., Tsuneyoshi, M. Intratumorous distribution of catecholaminergic clone cells of human neuroblastoma. A catecholamine fluorescence study. *Virchows Arch. [A]* 409:571-582, 1986.
37. Zaizen, Y., **Nakagawara, A.**, Ikeda, K. Patterns of destruction of mouse neuroblastoma cells by extracellular hydrogen peroxide formed by 6-hydroxydopamine and ascorbate. *J. Cancer Res. Clin. Oncol.* 111:93-97, 1986.
38. **Nakagawara, A.**, Ikeda, K., Tsuda, T., Higashi, K., Okabe, T. Amplification of N-myc oncogene in stage II and IVS neuroblastomas may be a prognostic indicator. *J. Pediatr. Surg.*, 22:415-418, 1987.
39. Tsuda, T., Obara, M., Hirano, H., Gotoh, S., Kubomura, S., Higashi, K., Kuroiwa, A., **Nakagawara, A.**, Nagahara, N., Shimizu, K. Analysis of N-myc amplification in relation to disease stage and histologic types in human neuroblastomas. *Cancer*, 60:820-826, 1987.
40. **Nakagawara, A.**, Ikeda, K. N-myc oncogene amplification and catecholamine metabolism in children with neuroblastoma. *Lancet*, 1:559, 1987.
41. **Nakagawara, A.**, Ikeda, K., Tsuda, T., Higashi, K. N-myc oncogene amplification and prognostic factors of neuroblastoma in children. *J. Pediatr. Surg.*, 22:895-898, 1987.
42. Hachitanda, Y., Tsuneyoshi, M., Daimaru, Y., Enjoji, M., **Nakagawara, A.**, Ikeda, K., Sueishi, K. Extraskeletal myxoid chondrosarcoma in young children. *Cancer* 61:2521-2526, 1988.
43. Iwanaga, M., **Nakagawara, A.**, Matsuo, S., Ikeda, K. Impaired polymorphonuclear leukocyte function in biliary atresia: role of bilirubin and bile acids. *J. Pediatr. Surg.* 22:967-972, 1987.
44. Sue, K., **Nakagawara, A.**, Okuzono, S., Fukushige, T., Ikeda, K. Combined effects of vitamin E (alpha-tocopherol) and cisplatin on the growth of murine neuroblastoma in vivo. *Eur. J. Cancer Clin. Oncol.* 24:1751-1758, 1988.
45. **Nakagawara, A.**, Ikeda, K., Tasaka, H. Dopaminergic neuroblastoma as a poor prognostic subgroup. *J. Pediatr. Surg.*, 23:346-349, 1988.
46. Hara, T., Mizuno, Y., Ishii, E., Ueda, K., Hirata, T., Daimaru, H., Enjoji, M., **Nakagawara, A.** Histiocytosis X presenting as multiple intrahepatic nodules. *Acta Jap. Hematol.* 51:1059-1062, 1988.

47. Sue, K., **Nakagawara, A.**, Okuzono, S., Fukushige, T., Ikeda, K. Different effects of vitamin E on the growth and morphological differentiation of 4 murine neuroblastoma cell lines in vitro. *Z. Kinderchir.* 43:398-401, 1988.
48. Okuzono, S., **Nakagawara, A.**, Sue, K., Fukushige, T., Ikeda, K. Different clonal drug sensitivity of murine neuroblastoma cells in vivo. *J. Pediatr. Surg.* 23:962-966, 1988.
49. **Nakagawara, A.**, Hachitanda, Y., Izbicki, T., Ikeda, K. Positive S100 protein staining of neuroblastoma discovered by VMA mass screening [letter]. *Eur. J. Pediatr.* 148:171-172. 1988.
50. **Nakagawara, A.**, Ikeda, K., Yokoyama, T., Tsuda, T., Higashi, K. Surgical aspects of N-myc oncogene amplification of neuroblastoma. *Surgery* 104:34-40, 1988.
51. **Nakagawara, A.**, Ikeda, K., Tsuda, T., Higashi, K. Biological characteristics of N-myc amplified neuroblastoma in patients over one year of age. *Progr. Clin. Biol. Res.* 271: 31-39, 1988.
52. Hashimoto, H., Daimaru, Y., Enjoji, M., **Nakagawara, A.** N-myc gene product expression in neuroblastoma. *J. Clin. Pathol.* 42: 52-55, 1989.
53. Ikeda, K., **Nakagawara, A.**, Yano, H., Akiyama, H., Tasaka, H., Ueda, K., Hara, T., Ishii, E., Ohgami, H., Sera, Y., Shimoda, H., Kishida, K., Tobo, Y., Hiyoshi, Y., Kawakami, K., Ohmori, H., Kuwano, S. Improved survival rates in children over 1 year of age with stage III or IV neuroblastoma following an intensive chemotherapeutic regimen. *J. Pediatr. Surg.* 24:189-193, 1989.
54. Fukushige, T., Ikeda, K., **Nakagawara, A.**, Nagoshi, M., Hashitanda, Y., Enjoji, M. Establishment of a cell line derived from congenital soft tissue tumor (primitive neuroectodermal tumor) in the right temporofacial region. *Human Cell* 2:208-209, 1989.
55. Ikeda, K., **Nakagawara, A.** Results of combination chemotherapy for advanced neuroblastoma. Kyushu Neuroblastoma Study Group. *Pediatr. Oncol.* 25:81-86, 1989.
56. Hachitanda, Y., **Nakagawara, A.**, Ikeda, K. An unusual anterior chest wall tumor due to actinomycosis in a child. *Pediatr. Radiol.* 20:96,1989.
57. Hachitanda, Y., Tsuneyoshi, M., Ehjoji, M., **Nakagawara, A.**, Shimada, H. Organoid neuroblastoma, a newly described entity. *Pediatr. Pathol.* 9:796, 1989.
58. Sue, K., Ikeda, K., **Nakagawara, A.**, Hironaga, Y., Hachitanda, Y., Otubo, K., Mitsuishi, T., Aoyama, T. Intrahepatic arterial injections of cisplatin-phosphatidylcholine- Lipiodol suspension in two unresectable hepatoblastoma cases. *Med. Pediatr. Oncol.* 17:496-500, 1989
59. Kawana, T., Ikeda, K., **Nakagawara, A.**, Kajiwara, M., Fukazawa, M., Hara, K. A case of VACTEL syndrome with antenatally diagnosed duodenal atresia. *J. Pediatr. Surg.* 24:1158-1160, 1989.
60. **Nakagawara, A.**, Tsuda, T., Higashi, K. Amplification of clone 8 as a prognostic indicator of neuroblastoma [letter]. *Eur. J. Pediatr.* 148:684, 1989.
61. **Nakagawara, A.** The persistent challenge of neuroblastoma: Role of N-myc oncogene and prognosis. In: Kimura, K., Ota, K., Carter, S.K., Pinedo, H.M. (eds): *Cancer Chemotherapy: Challenges for the Future*. Excerpta Medica, Tokyo, 1989, pp. 208-213.
62. Nagao, M., Ishizaka, Y., **Nakagawara, A.**, Kohno, K., Kuwano, M., Tahira, T., Itoh, F., Ikeda, I., Sugimura, T. Expression of ret proto-oncogene in human neuroblastomas. *Jpn. J. Cancer Res.*, 81:309-312, 1990.

63. **Nakagawara, A.**, Ikeda, K., Higashi, K., Sasazuki, T. Inverse correlation between N-myc amplification and catecholamine metabolism in children with advanced neuroblastoma. *Surgery*, 107:43-49, 1990.
64. **Nakagawara, A.**, Kadomatsu, K., Sato, S., Kohno, K., Takano, H., Akazawa, K., Nose, Y., Kuwano, M. Inverse correlation between expression of multidrug resistance gene and N-myc oncogene in human neuroblastomas. *Cancer Res.* 50: 3043-3047, 1990.
65. **Nakagawara, A.**, Sasazuki, T., Akiyama, H., Kawakami, K., Kuwano, A., Yokoyama, T., Kume, K. N-myc oncogene and stage IV-S neuroblastoma. Preliminary observations in 10 cases. *Cancer* 65:1960-1967, 1990.
66. Sano, K., Nakamura, H., Mabuchi, S., Tanaka, T., **Nakagawara, A.**, Takai, Y. Expression of the smg p25A (a ras p21-like GTP-binding protein) gene in human neuroblastoma cell lines and tumor tissues. *Cancer Res.* 50:7242-7245, 1990.
67. Hachitanda, Y., Tsuneyoshi, M., Enjoji, M., **Nakagawara, A.**, Ikeda, K. Congenital primitive neuroectodermal tumor with epithelial and glial differentiation. An ultrastructural and immunohistochemical study. *Arch Pathol Lab Med* 114:101-105, 1990.
68. Sugio, K., **Nakagawara, A.**, Sasazuki, T. Association of expression between N-myc gene and major histocompatibility complex class I gene in surgically resected human neuroblastoma. *Cancer* 67:1384-1388, 1991
69. **Nakagawara, A.**, Zaizen, Y., Ikeda, K., Suita, S., Ohgami, H., Nagahara, N., Sera, Y., Akiyama, H., Kawakami, K., Uchino, J.-i. Different genomic and metabolic patterns between mass screening-positive and mass screening-negative later-presenting neuroblastomas. *Cancer* 68:2037-2044, 1991.
70. **Nakagawara, A.**, Okabe, I., Morita, K., Uchino, J., Ohi, R., Iwafuchi, M., Matsuyama, S., Nagashima, K., Takahashi, H., Nakajo, T., Hirai, Y., Tsuchida, Y., Saeki, M., Yokoyama, J., Nishi, T., Okamoto, E., Suita, S. Proposal and assessment of Japanese Tumor Node Metastasis postsurgical histopathological staging system for neuroblastoma based on an analysis of 495 cases. *Jpn. J. Clin. Oncol.* 21:1-7, 1991
71. **Nakagawara, A.**, Kadomatsu, K., Sato, S., Kohno, K., Takano, H., Kuwano, M. Inverse expression of MYCN and mdr-1 in human neuroblastoma. *Progr. Clin. Biol. Res.* 366: 11-19, 1991.
72. **Nakagawara, A.**, Zaizen, Y., Suita, S. Genomic and metabolic aspects of neuroblastomas in children under one year of age. *Progr. Clin. Biol. Res.* 366: 115-125, 1991.
73. Fukushige, T., Nagoshi, M., Hachitanda, Y., Ueno, T., Zaizen, Y., Suita, S., Tsuneyoshi, M., Kaneko, Y., **Nakagawara, A.** Establishment and characterization of a cell line of congenital primitive neuroectodermal tumor of soft tissue. *Virchows Archiv [B Cell Pathol]* 62:159-166, 1992
74. Hirotsune, S., Hatada, I., Komatsubara, H., Nagai, H., Kuma, K., Kobayakawa, K., Kawara, T., **Nakagawara, A.**, Fujii, K., Mukai, T., Hayashizaki, Y. New approach for detection of amplification in cancer DNA using restriction landmark genomic scanning. *Cancer Res* 52:3642-3647, 1992
75. Yamada, S., Ishii, E., **Nakagawara, A.**, Akazawa, K., Tasaka, H., Kajiwara, M., Kawakami, K., Ueda, K. Features and outcome of advanced neuroblastoma with distant lymph node metastasis. *Pediatr. Hematol. Oncol.* 9:49-56, 1992

76. Kadomatsu, K., **Nakagawara, A.**, Zaizen, Y., Nagoshi, M., Tsuneyoshi, M., Fukushige, T., Suita, S. Undifferentiated (embryonal) sarcoma of the liver: Report of three cases. *Jpn. J. Surg.* 22:451-455, 1992.
77. **Nakagawara, A.**, Arima, M., Azar, C.G., Scavarda, N.J., Brodeur, G.M. Inverse relationship between *trk* expression and *Nmyc* amplification in human neuroblastomas. *Cancer Res.* 52:1364-1368, 1992.
78. Maminta, M.L.D., Williams, K.W., **Nakagawara, A.**, Enger, K.T., Guo, C., Brodeur, G.M., Deuel, T.F. Identification of a novel tyrosine kinase receptor-like molecule in neuroblastomas. *Biochem. Biophys. Res. Commun.* 189:1077-1083, 1992.
79. Hachitanda, Y., **Nakagawara, A.**, Nagoshi, M., Tsuneyoshi, M. Prognostic value of N-myc oncogene amplification and S-100 protein positivity in children with neuroblastic tumors. *Acta Pathol. Jpn.* 42:639-644, 1992.
80. Murakami, T., Ohmori, H., Katoh, T., Nakamura, H., Tsuda, T., Shan, R. C., Akiya, S., **Nakagawara, A.**, Higashi, K. Differences of L-myc polymorphic patterns of neuroblastoma in patients under 1 year versus older ages: A preliminary report. *J. Pediatr. Surg.* 27:1442-1446, 1992.
81. Brodeur, G. M., **Nakagawara, A.** Molecular basis for clinical heterogeneity in neuroblastoma. *Am. J. Pediatr. Hematol. Oncol.* 14:111-116, 1992.
82. Brodeur, G. M., Azar, C., Brother, M., Hiemstra, J., Kaufman, B., Marshall, H., Moley, J., **Nakagawara, A.**, Saylor, R., Scavarda, N., Schneider, S., Wasson, J., White, P., Seeger, R., Look, T., Castleberry, R. Neuroblastoma: Effect of genetic factors on prognosis and treatment. *Cancer* 70: 1685-1694, 1992.
83. Tsutsui, J., Kadomatsu, K., Matsubara, S., **Nakagawara, A.**, Hamanoue, M., Takao, S., Shimazu, H., Ohi, Y., Muramatsu, T. A new family of heparin-binding growth/differentiation factors: Increased Midkine expression in Wilms' tumor and other human carcinomas. *Cancer Res.* 53:1281-1285, 1993.
84. **Nakagawara, A.**, Arima-Nakagawara, M., Scavarda, N.J., Azar, C.G., Cantor, A.B., Brodeur, G.M. Association between high levels of expression of the TRK gene and favorable outcome in human neuroblastoma. *N. Engl. J. Med.* 328:847-854, 1993.
85. Muraji, T., Fujimoto, J., Okamoto, E., Suita, S., **Nakagawara, A.** Combined determination of N-myc oncogene amplification and DNA ploidy in neuroblastoma: Complementary prognostic indicators. *Cancer* 72:2763-2768, 1993.
86. **Nakagawara, A.**, Azar, C.G., Scavarda, N.J., Brodeur, G.M. Expression and function of TRK-B and BDNF in human neuroblastomas. *Mol. Cell. Biol.* 14:759-767, 1994.
87. **Nakagawara, A.**, Arima-Nakagawara, M., Azar, C. G., Scavarda, N. J., Brodeur, G. M. Clinical significance of expression of neurotrophic factors and receptors in neuroblastoma. In A. E. Evans, J. L. Biedler, G. M. Brodeur, G. J. D'Angio, and A. Nakagawara (eds), *Advances in Neuroblastoma Research 4*, pp 155-161, Wiley-Liss, Inc.: New York, 1994.
88. Azar, C. G., Scavarda, N. J., **Nakagawara, A.**, Brodeur, G. M. Expression and function of the nerve growth factor receptor (TRK-A) in human neuroblastoma cell lines. In A. E. Evans, J. L. Biedler, G. M. Brodeur, G. J. D'Angio, and A. Nakagawara (eds), *Advances in Neuroblastoma Research 4*, pp 169-175, Wiley-Liss, Inc.: New York, 1994.
89. Brodeur, G. M. and **Nakagawara, A.** Molecular aspects of neuroblastoma and its prognosis. In T. Sawada, T. Matsumura, and Z. Kizaki (eds), *Proceedings of the 3rd International Symposium on Neuroblastoma Screening*, pp 159-164, 1994.

90. **Nakagawara, A.**, Liu, X., Ikegaki, N., White, P., Yamashiro, D. J., Nycum, L. M., Biegel, J. A., Brodeur, G. M. Cloning and chromosomal localization of human TRK-B (NTRK2) tyrosine kinase receptor gene. *Genomics* 25:538-546, 1995.
91. **Nakagawara, A.**, Milbrandt, J., Muramatsu, T., Deuel, T. F., Zhao, H., Cnaan, A., Brodeur, G. M. Differential expression of pleiotrophin and midkine in advanced neuroblastomas. *Cancer Res.* 55:1792-1797, 1995.
92. Ikegaki, N., Katsumata, M., Tsujimoto, Y., **Nakagawara, A.**, Brodeur, G. M. Expression of bcl-2 relative to myc expression in human neuroblastomas. *Cancer Lett.* 91:161-168, 1995.
93. Yamashiro, D., **Nakagawara, A.**, Ikegaki, N., Liu, X., Brodeur, G. M. Expression of TRK-C in favorable human neuroblastomas. *Oncogene* 12:37-41, 1996.
94. Reale, M.A., Reyes-Mugica, M., Pierceall, W.E., Rubinstein, M.C., Hedrick, L., Cohn, S.L., **Nakagawara, A.**, Brodeur, G.M., Fearon, E.R. Loss of DCC expression in neuroblastoma is associated with disease dissemination. *Clin. Cancer Res.* 2:1097-1102, 1996.
95. Ozaki, T., Hishiki, T., Toyama, Y., Yuasa, S., **Nakagawara, A.**, Sakiyama, S. Identification of a new cellular protein which can interact specifically with DAN. *DNA Cell Biol.* 16:985-991, 1996
96. Kimura, M., Tagawa, M., Takenaga, K., Yamaguchi, T., Saisho, H., **Nakagawara, A.**, Sakiyama, S. Inability to induce the alteration of tumorigenicity and chemosensitivity of p53-null human pancreatic carcinoma cells by the transduction of wild-type p53 gene. *Anticancer Res.* 17:879-884, 1997.
97. Kimura, M., Tagawa, M., Takenaga, K., **Nakagawara, A.**, Horitsu, K., Yamaguchi, T., Saisho, H., Sakiyama, S. Drug resistance to ganciclovir observed in suicide gene therapy is due to the loss of integrated herpes simplex virus-thymidine kinase gene. *Intl. J. Oncol.* 10:775-778, 1997.
98. **Nakagawara, A.**, Nakamura, Y., Ikeda, H., Hiwasa, T., Kuida, K., Su, M. S.-S., Zhao, H., Cnaan, A., Sakiyama, S. High levels of expression and nuclear localization of ICE and CPP32 in favorable human neuroblastomas. *Cancer Res.* 57:4578-4584, 1997.
99. Ozaki, T., Enomoto, H., Nakamura, Y., Kondo, K., Seki, N., Ohira, M., Nomura, N., Ohki, M., **Nakagawara, A.**, Sakiyama, S. The genomic analysis of human DAN gene. *DNA Cell Biology* 16:1031-1039, 1997.
100. Ozaki, T., Kondo, K., Nakamura, Y., Ichimiya, S., **Nakagawara, A.**, Sakiyama, S. Interaction of DA41, a DAN-binding protein, with the epidermal growth factor-like protein, S(1-5). *Biochem. Biophys. Res. Commun.* 237:245-250, 1997.
101. Kageyama, H., Seki, N., Yamada, S., Sakiyama, S., **Nakagawara, A.** DPC4 splice variants in neuroblastoma. *Cancer Lett.* 122:187-193, 1997.
102. Seki, N., Nimura, Y., Ohira, M., Saito, T., Ichimiya, S., Nomura, N., **Nakagawara, A.** Identification and chromosome assignment of a human gene encoding a novel phosphatidylinositol-3 kinase. *DNA Research* 4:355-358, 1997.
103. Hiwasa, T., Kondo, K., Hishiki, T., Koshizawa, S., Umezawa, K., **Nakagawara, A.** GDNF-induced neurite formation was stimulated by protein kinase inhibitors and suppressed by Ras inhibitors. *Neurosci. Lett.* 238:115-118, 1997.
104. Brodeur, G. M., **Nakagawara, A.**, Yamashiro, D. J., Ikegaki, N., Liu, X-G., Azar, C. G., Lee, C. P., Evans, A. E. Expression of TrkA, TrkB and TrkC in human neuroblastomas. *J. Neuro-Oncol.* 31:49-55, 1997.

105. **Nakagawara, A.**, Brodeur, G. M. Role of neurotrophins and their receptors in human neuroblastomas: A primary culture study. *Eur. J. Cancer* 33:2050-2053, 1997.
106. Ikeda, H., Nakamura Y., Hiwasa, T., Sakiyama, S., Kuida, K., Su, M. S-S., **Nakagawara, A.** Interleukin-1 b converting enzyme (ICE) is preferentially expressed in neuroblastomas with favorable prognosis. *Eur. J. Cancer* 33:2081-2083, 1997.
107. Nakamura, Y., Ozaki, T., Enomoto, H., Kondo, K., **Nakagawara, A.**, Sakiyama, S. A product of DAN, a novel candidate tumor suppressor gene, is secreted into culture medium and suppresses DNA synthesis. *Eur. J. Cancer* 33:1986-1990, 1997.
108. Yamashiro, D. J., Liu, X., Lee, C., **Nakagawara, A.**, Ikegaki, N., McGregor, L. M., Baylin, S. B., Brodeur, G. M. Expression and function of TrkC in human neuroblastoma. *Eur. J. Cancer* 33:2054-2057, 1997.
109. Hiwasa, T., Tokita, H., Sakiyama, S. and **Nakagawara, A.** Enhancement of chemosensitivity towards anticancer drugs by high expression of caspase-1 in NIH3T3 cells. *Anti-cancer Drugs* 9:82-87, 1998.
110. Hishiki, T., Nimura, Y., Isogai, E., Kondo, K., Ichimiya, S., Nakamura, Y., Ozaki, T., Sakiyama, S., Hirose, M., Seki, N., Takahashi, H., Ohnuma, N., Tanabe, M., **Nakagawara, A.** Glial cell line-derived neurotrophic factor/neurturin-induced differentiation and its enhancement by retinoic acid in primary human neuroblastomas expressing c-Ret, GFR α -1 and GFR α -2. *Cancer Res.* 58:2158-2165, 1998.
111. Takahashi, H., Ichimiya, S., Watanabe, M., Furusato, M., Wakui, S., Yatani, R., Aizawa, S., **Nakagawara, A.** Mutation, allelotyping, and transcription analyses of the p73 gene in prostatic carcinoma. *Cancer Res.* 58:2076-2077, 1998.
112. Yoshida, H., Enomoto, H., Tagawa, M., Takenaga, K., Tasaki, K., **Nakagawara, A.**, Ohnuma, N., Takahashi, H., Sakiyama, S. Impaired tumorigenicity and decreased liver metastasis of murine neuroblastoma cells engineered to secrete interleukin-2 or granulocyte macrophage-colony stimulating factor. *Cancer Gene Therapy* 5:67-73, 1998.
113. Hiwasa, T., Kondo, K., **Nakagawara, A.**, Ohkoshi, M. Potent growth-suppressive activity of a serine protease inhibitor, ONO-3403, toward malignant human neuroblastoma cell lines. *Cancer Lett.* 126:221-225, 1998.
114. Nakamura, Y., Ozaki, T., Ichimiya, S., **Nakagawara, A.**, Sakiyama, S. Ectopic expression of DAN enhances the retinoic acid-induced neuronal differentiation in human neuroblastoma cell lines. *Biochem. Biophys. Res. Commun.* 243:722-726, 1998.
115. Nishio, S., Morioka, T., Hamada, Y., Fukui, M., **Nakagawara, A.** Immunohistochemical expression of tyrosine kinase (Trk) receptor proteins in mature neuronal cell tumors of the central nervous system. *Clin. Neuropathol.* 17:123-130, 1998.
116. Mori, K., Iwao, K., Miyoshi, Y., **Nakagawara, A.**, Kofu, K., Akiyama, T., Arita, N., Hayakawa, T., Nakamura, Y. Identification of brain-specific splicing variants of the hDLG1 gene and altered splicing in neuroblastoma cell lines. *J. Hum. Genet.* 43:123-127, 1998.
117. **Nakagawara, A.** The NGF Story and Neuroblastoma. *Med. Pediatr. Oncol.* 31:113-115, 1998.
118. Enomoto, H., Nakamura, Y., Kondo, K., Hirose, M., Kaneko, Y., Horie, H., Takahashi, H., Ohnuma, N., Tanabe, M., **Nakagawara, A.** In vivo karyotypic evolution with altered biological characteristics in a case of neuroblastoma. *J. Pediatr. Surg.* 33:1404-1407, 1998.

119. Kimura, M., Tagawa, M., Takenaga, K., Kondo, F., Yamaguchi T., Saisho H., **Nakagawara, A.**, Sakiyama, A. Loss of tumorigenicity of human pancreatic carcinoma cells engineered to produce IL-2 and IL-4 in nude mice: a potentiality for cancer gene therapy. *Cancer Lett.* 128:47-53, 1998.
120. Osada, M., Ohba, M., Kawahara, C., Ishioka, C., Kanamaru, R., Katoh I., Ikawa, Y., Nimura, Y., **Nakagawara, A.**, Obinata, M., Ikawa, S. Cloning and functional analysis of human p51, structurally and functionally resembling p53. *Nat. Med.* 4:839-843, 1998.
121. Sunahara, M., Ichimiya, S., Nimura, Y., Takada, N., Sakiyama, S., Sato, Y., Todo, S., Adachi, W., Amano, J., **Nakagawara, A.** Mutational analysis of the p73 gene localized at chromosome 1p36.3 in colorectal carcinomas. *Int. J. Oncol.* 13:319-323, 1998.
122. Seki, N., Sugano, S., Suzuki, Y., **Nakagawara, A.**, Ohira, M., Muramatsu, M., Saito, T., Hori T. Isolation, tissue expression and chromosomal assignment of human RGS5, a novel G-protein signaling regulator gene. *J. Hum. Genet.* 43:202-205, 1998.
123. Uchiyama, K., Muramatsu, H., Kaname, T., Ogawa, H., Yamakawa, T., Fan, Q-W., Mitsuoka, C., Kannagi, R., Habuchi, O., Yokoyama, I., Yamamura, K., Ozaki, T., **Nakagawara, A.**, Kadomatsu K., Muramatsu, T. Human N-acetylglucosamine-6-O-sulfotransferase involved in the biosynthesis of 6-sulfo sialyl Lewis X: Molecular cloning, chromosomal mapping, and expression in various organs and tumor cells. *J. Biochem.* 124:670-678, 1998.
124. Nimura, Y., Mihara, M., Ichimiya, S., Sakiyama, S., Seki, N., Ohira, M., Nomura, N., Fujimori, M., Adachi, W., Amano, J., He, M., Ping, Y-M., **Nakagawara, A.** p73, a gene related to p53, is not mutated in esophageal carcinomas. *Int. J. Cancer* 78:437-440, 1998.
125. Hiwasa, T., **Nakagawara, A.** Suppression of transformed phenotypes of Ha-ras-transformed NIH3T3 cells by caspase-2. *Biochem. Biophys. Res. Commun.* 250:741-746, 1998.
126. He, L., Yu, J. X., Liu, L., Buyse, I. M., Wang, M., Yang, Q., **Nakagawara, A.**, Brodeur, G. M., Shi, Y. E., Huang, S. RIZ1, but not the alternative RIZ2 product of the same gene, is underexpressed in breast cancer, and forced RIZ1 expression causes G2-M cell cycle arrest and/or apoptosis. *Cancer Res.* 58:4238-4244, 1998.
127. Seki, N., Muramatsu, M., Sugano, S., Suzuki, Y., **Nakagawara, A.**, Ohira, M., Hayashi, A., Hori, T., Saito, T. Cloning, expression analysis, and chromosomal localization of HIP1R, an isolog of Huntingtin interacting protein (HIP1). *J. Hum. Genet.* 43:268-271, 1998.
128. Seki, N., Hattori, A., Sugano, S., Suzuki, Y., **Nakagawara, A.**, Ohira, M., Muramatsu, M., Hori, T., Saito, T. Isolation, tissue expression, and chromosomal assignment of a novel human gene which encodes a protein with RING finger motif. *J. Hum. Genet.* 43:272-274, 1998.
129. **Nakagawara, A.** Molecular basis of spontaneous regression of neuroblastoma: Role of neurotrophic signals and genetic abnormalities. *Human Cell* 11:115-124, 1998.
130. Hiwasa, T., Nakamura, Y., Ozaki, T., Kondo, K., Saido, T., **Nakagawara, A.**, Sakiyama, S. Down-regulation of protein kinase C α and γ and enhanced TPA-induced neurite formation in DAN-transfected neuroblastoma cells. *FEBS Lett.* 440:25-28, 1998.
131. Kaneko, Y., Kobayashi, H., Maseki, N., **Nakagawara, A.**, Sakurai, M. Disomy 1 with terminal 1p deletion was frequent in mass screening-negative/late presenting neuroblastomas in young children, but not in mass screening-positive neuroblastomas in infants. *Int. J. Cancer* 80:54-59, 1999.
132. Mihara, M., Nimura, Y., Ichimiya, S., Sakiyama, S., Kajikawa, S., Adachi, W., Amano, J., **Nakagawara, A.** Absence of mutation of the p73 gene localized at chromosome 1p36.3 in hepatocellular carcinoma. *Br. J. Cancer* 79:164-167, 1999.

133. Ichimiya, S., Nimura, Y., Kageyama, H., Takada, N., Sunahara, M., Shishikura, T., Nakamura Y., Sakiyama, S., Seki, N., Ohira, M., Kaneko, Y., McKeon, F., Caput, D., **Nakagawara, A.** p73 at chromosome 1p36.3 is lost in advanced stage neuroblastoma but its mutation is infrequent. *Oncogene* 18:1061-1066, 1999.
134. Seki, N., Hattori, A., Sugano, S., Suzuki, Y., **Nakagawara, A.**, Muramatsu, M., Hori, T., Saito, T. A novel human gene whose product shares significant homology with the bovine brain-specific protein p25 on chromosome 5p15.3. *J. Hum. Genet.* 44:121-122, 1999.
135. Yang, H-W., Ikeda, H., Kato, K., **Nakagawara, A.**, Choi, S. H., Hayashi, Y., Hemmi, H., Tsuchida, Y. Development of a polyclonal antibody with defined specificity against synthetic peptides from the N-myc oncoprotein, using multiple antigen peptide and hemocyanin conjugation methods. *J. Pediatr. Surg.* 34:454-460, 1999.
136. Han, S., Semba, S., Abe, T., Makino, N., Furukawa, T., Fukushige, S., Takahashi, H., Sakurada, A., Sato, M., Shiiba, K., Matsuno, S., Nimura, Y., **Nakagawara, A.**, Horii, A. Infrequent somatic mutations of the p73 gene in various human cancers. *Eur. J. Surg. Oncol.* 25:194-198, 1999.
137. Nishio, S., Hamada, Y., **Nakagawara, A.**, Haga, S., Suzuki, S., Fukui, M. Thoracic paravertebral ganglioneuroma with high immunohistochemical expression of TrkA. *Neuropathol.* 19:51-56, 1999.
138. Shishikura, T., Ichimiya, S., Ozaki, T., Nimura, Y., Kageyama, H., Nakamura, Y., Sakiyama, S., Miyauchi, M., Yamamoto, N., Suzuki, M., Nakajima, N., **Nakagawara, A.** Mutational analysis of the p73 gene in human breast cancers. *Int. J. Cancer* 84:321-325, 1999.
139. Takada, N., Ozaki, T., Ichimiya, S., Todo, S., **Nakagawara, A.** Identification of a transactivation activity in the carboxy-terminal region of p73 which is impaired in the naturally occurring mutants found in human neuroblastomas. *Cancer Res.* 59:2810-2814, 1999.
140. Hirai, M., Yoshida, S., Kashiwagi, H., Kawamura, T., Ishikawa, T., Kaneko, M., Ohkawa, H., **Nakagawara, A.**, Miwa, M., Uchida, K. 1q23 gain is associated with progressive neuroblastoma resistant to aggressive treatment. *Genes Chromosomes Cancer* 25:261-269, 1999.
141. Ozaki, T., Hanaoka, E., Naka, M., **Nakagawara, A.**, Sakiyama, S. Cloning and characterization of rat *BAT3* cDNA. *DNA Cell Biol.* 18:503-512, 1999.
142. Sunahara, M., Shishikura, T., Takahashi, M., Todo, S., Yamamoto, N., Kimura, H., Kato, S., Ishioka, C., Ikawa, S., Ikawa, Y., **Nakagawara, A.** Mutational analysis of p51A/TAp63 γ , a p53 homolog, in non-small cell lung cancer and breast cancer. *Oncogene* 18:3761-3765, 1999.
143. Ozaki, T., Naka, M., Takada, N., Tada, M., Sakiyama, S., **Nakagawara, A.** Deletion of the COOH-terminal region of p73 α enhances both its transactivation function and DNA-binding activity but inhibits induction of apoptosis in mammalian cells. *Cancer Res.* 59:5902-5907, 1999.
144. Kato, S., Shimada, A., Osada, M., Ikawa, S., Obinata, M., **Nakagawara, A.**, Kanamaru, R., Ishioka, C. Effects of p51/p63 missense mutations on transcriptional activities of p53 downstream gene promoters. *Cancer Res.* 59:5908-5911, 1999.
145. Ikawa S., **Nakagawara A.**, Ikawa Y. p53 family genes: structural comparison, expression and mutation. *Cell Death Differ.* 6:1154-1161, 1999.
146. Sugimoto, N., Fukuda, Y., Saito-Ohara, F., Kamiyama, R., **Nakagawara, A.**, Mukae, N., Nagata, S., Inazawa, J. The human caspase-activated DNase (CAD) gene: genomic structure, exonic single-nucleotide polymorphisms, and a highly polymorphic dinucleotide repeat at the CAD locus. *J. Hum. Genet.* 44:408-411, 1999.

147. Islam, A., Kageyama, H., Takada, N., Kawamoto, T., Takayasu, H., Isogai, E., Ohira, M., Hashizume, K., Kobayashi, Y., Kaneko, Y., **Nakagawara, A.** High expression of *Survivin*, mapped to 17q25, is significantly associated with poor prognostic factors and promotes cell survival in human neuroblastoma. *Oncogene* 19:617-623, 2000.
148. Nagai, M., Ichimiya, S., Ozaki, T., Seki, N., Mihara, M., Furuta, S., Ohira, M., Tomioka, N., Nomura, N., Sakiyama, S., Kubo, O., Takakura, K., Hori, T., **Nakagawara, A.** Identification of the full-length KIAA0591 gene encoding a novel kinesin-related protein which is mapped to the neuroblastoma suppressor gene locus at 1p36.2. *Int. J. Oncol.* 16:907-916, 2000.
149. Hanaoka E, Ozaki T, Ohira M, Nakamura Y, Suzuki M, Takahashi E, Moriya H, **Nakagawara A**, Sakiyama S. Molecular cloning and expression analysis of the human DA41 gene and its mapping to chromosome 9q21.2-q21.3. *J. Hum. Genet.* 45:188-191, 2000.
150. **Nakagawara A**, Kogner P. Expression and function of Trk and its related genes in human neuroblastoma. In Neuroblastoma, edited by Brodeur GM, Sawada T, Tsuchida Y, Voute PA, Elsevier, Amsterdam, pp147-157, 2000.
151. Igarashi, J., Nimura, Y., Fujimori, M., Mihara, M., Adachi, W., Kageyama, H., **Nakagawara, A.** Allelic loss of the region of chromosome 1p35-pter is associated with progression of human gastric carcinoma. *Jpn. J. Cancer Res.* 91:797-801, 2000.
152. Tsujimoto T, Mochizuki S, Iwadata Y, Namba H, Nagai M, Kawamoto T, Sunahara M, Yamaura A, **Nakagawara A**, Sakiyama S, Tagawa M. The p73 gene is not mutated in oligodendroglioma which frequently have a deleted region at chromosome 1p36.3. *Anti-Cancer Res.* 20:2495-2498, 2000.
153. Ozaki, T., Nagase, T., Ichimiya, S., Seki, N., Ohira, M., Nomura, N., Takada, N., Sakiyama, S., Weber, B.L., **Nakagawara, A.** NFBD1/KIAA0170 is a novel nuclear transcriptional transactivator with BRCT domain. *DNA Cell Biol.* 19:475-485, 2000.
154. Ohira M, Kageyama H, Mihara M, Furuta S, Machida T, Shishikura T, Takayasu H, Islam A, Nakamura Y, Takahashi M, Tomioka N, Sakiyama S, Kaneko Y, Toyoda A, Hattori M, Sakaki Y, Ohki M, Horii A, Soeda E, Inazawa J, Seki N, Kuma H, Nozawa I, **Nakagawara A** Identification and characterization of a 500-kb homozygously deleted region at 1p36.2-p36.3 in a neuroblastoma cell line. *Oncogene* 19:4302-4307, 2000.
155. Ichimiya S, **Nakagawara A**, Sakuma Y, Kimura S, Ikeda T, Satoh M, Takahashi N, Sato N, Mori M. p73: Structure and function. *Pathol. Internatl.* 50:589-593, 2000.
156. Ozaki T, Nakamura Y, Hanaoka E, **Nakagawara A**, Sakiyama S. Overexpression of DA41 in v-Ha-ras-3Y1 cells causes a growth suppression associated with the accumulation of p21^{WAF1/CIP1}. *Jpn. J. Cancer Res.* 91:1-7, 2000.
157. Hanaoka E, Ozaki T, Nakamura Y, Moriya H, **Nakagawara A**, Sakiyama S. Overexpression of DAN causes a growth suppression in p53-deficient SAOS-2 cells. *Biophys. Biochem. Res. Commun.* 278:20-26, 2000.
158. Ohira M, Shishikura T, Kawamoto T, Inuzuka H, Morohashi A, Takayasu H, Kageyama H, Takada N, Takahashi M, Sakiyama S, Suzuki Y, Sugano S, Kuma H, Nozawa I, **Nakagawara A.** Hunting the subset-specific genes of neuroblastoma: expression profiling and differential screening of the full-length-enriched oligo-capping cDNA libraries. *Med. Pediatr. Oncol.* 35:547-549, 2000.
159. Furuta S, Ohira M, Machida T, Hamano S, **Nakagawara A.** Analysis of loss of heterozygosity at 16p12-p13 (familial neuroblastoma locus) in 470 neuroblastomas including both sporadic and mass screening tumors. *Med. Pediatr. Oncol.* 35:531-533, 2000.

160. **Nakagawara A**, Ohira M, Kageyama H, Mihara M, Furuta S, Machida T, Takayasu H, Islam A, Nakamura Y, Takahashi M, Shishikura T, Kaneko Y, Toyoda A, Hattori M, Sakaki Y, Ohki M, Horii A, Soeda E, Inazawa J, Seki N, Kuma H, Nozawa I, Sakiyama S. Identification of the homozygously deleted region at chromosome 1p36.2 in human neuroblastoma. *Med. Pediatr. Oncol.* 35:516-521, 2000.
161. Kawamoto T, Shishikura T, Ohira M, Takayasu H, Morohashi A, Takada N, Takahashi M, Suzuki Y, Sugano S, Hori T, **Nakagawara A**. Association between favorable neuroblastoma and high expression of the novel metalloproteinase gene, nbla3145/XCE, cloned by differential screening of the full-length-enriched oligo-capping neuroblastoma cDNA libraries. *Med. Pediatr. Oncol.* 35:628-631, 2000.
162. Islam A, Kageyama H, Hashizume K, Kaneko Y, **Nakagawara A**. Role of survivin, whose gene is mapped to 17q25, in human neuroblastoma and identification of a novel dominant-negative isoform, survivin- β /2B. *Med. Pediatr. Oncol.* 35:550-553, 2000.
163. Yamauchi K, Tozuka M, Hidaka H, Nakabayashi T, Sugano M, Kondo Y, **Nakagawara A**, Katsuyama T. Effect of apolipoprotein AII on the interaction of apolipoprotein E with beta-amyloid: some apo(E-AII) complexes inhibit the internalization of beta-amyloid in cultures of neuroblastoma cells. *J. Neurosci. Res.* 62:608-614, 2000.
164. Ichimiya, S., Nimura, Y., Kageyama, H., Takada, N., Sunahara, M., Shishikura, T., Nakamura, Y., Sakiyama, S., Seki, N., Ohira, M., Kaneko, Y., McKeon, F., Caput, D., **Nakagawara, A**. Genetic analysis of p73 localized at chromosome 1p36.3 in primary neuroblastomas. *Med. Pediatr. Oncol.* 36:42-44, 2001.
165. Takada, N., Isogai, E., Kawamoto, T., Nakanishi, H., Todo, S., **Nakagawara, A**. Retinoic acid-induced apoptosis of CHP134 neuroblastoma cell line is associated with nuclear accumulation of p53 and is rescued by GDNF/Ret signal. *Med. Pediatr. Oncol.* 36:122-126, 2001.
166. Ichimiya, S., Nimura, Y., Seki, N., Ozaki, T., Nagase, T., **Nakagawara, A**. Down-regulation of hASH1 is associated with the retinoic acid-induced differentiation of human neuroblastoma cell lines. *Med. Pediatr. Oncol.* 36:132-134, 2001.
167. Kawamura K, Oh K, Bahar R, Koshikawa N, Shishikura T, **Nakagawara A**, Sakiyama S, Kajiwara K, Kimura M, Tagawa M. The error-prone DNA polymerase β catalytic subunit (Rev3) gene is ubiquitously expressed in normal and malignant human tissues. *Int. J. Oncol.* 18:97-103, 2001.
168. Hirai M, Takenaga K, Koshikawa N, **Nakagawara A**, Takenouchi T, Araki A, Oosaki T, Hayashi Y, Hirono M, Yoshihara T. Expression of vascular endothelial growth factors (VEGF-A/VEGF-1 and VEGF-C/VEGF-2) in postmenopausal uterine endometrial carcinoma. *Gynecol. Oncol.* 80:181-188, 2001.
169. Miyauchi, M., Yoshida, Y., Tada, Y., Narita, M., Maeda, T., Bahar, R., Kadomatsu, K., Muramatsu, T., Matsubara, S., **Nakagawara, A.**, Sakiyama, S., Tagawa, M. Expression of herpes simplex virus-thymidine kinase gene controlled by a promoter region of the midkine gene confers selective cytotoxicity to ganciclovir in human carcinoma cells. *Int. J. Cancer* 91:723-727, 2001.
170. Aoyama M, Asai K, Shishikura T, Kawamoto T, Miyauchi T, Yokoi T, Togari H, Wada Y, Kato T, **Nakagawara A**. Human neuroblastomas with unfavorable biologies express high levels of brain-derived neurotrophic factor mRNA and a variety of its variants. *Cancer Lett.* 164:51-60, 2001.
171. Takayasu H, Horie H, Hiyama E, Matsunaga T, Hayashi Y, Watanabe Y, Suita S, Kaneko M, Sasaki F, Hashizume K, Ozaki T, Furuuchi K, Tada M, Ohnuma N, **Nakagawara A**. Frequent

- deletions and mutations of the b-catenin gene are associated with overexpression of cyclin D1 and fibronectin and poorly-differentiated histology in childhood hepatoblastoma. *Clin.Cancer Res.* 7:901-908, 2001.
172. Ozaki T, **Nakagawara A.** p73 and p53: Functional implications of their interaction. *Recent Research Development in Cancer* 3:89-110, 2001.
173. Naka M, Ozaki T, Takada N, Takahashi M, Shishikura T, Sakiyama S, Tada M, Todo S, **Nakagawara A.** Functional characterization of naturally occurring mutants (P405R and P425L) of p73 α and p73 β found in neuroblastoma and lung cancer. *Oncogene* 20:3568-3572, 2001.
174. Chen YZ, Hayashi Y, Wu JG, Yakaoka E, Maekawa K, Watanabe N, Inazawa J, Hosoda F, Arai Y, Mizushima H, Morohashi A, Ohira M, **Nakagawara A,** Liu S-Y, Hosi M, Horii A, Soeda E. A BAC-based STS-content map spanning a 35-megabase region of human chromosome 1p35-36. *Genomics* 74:55-70, 2001.
175. **Nakagawara A.** Trk receptor tyrosine kinases: A bridge between cancer and neural development. *Cancer Lett.* 169:107-114, 2001.
176. Momoi H, Okabe H, Kamikawa T, Satoh S, Ikai I, Yamamoto M, **Nakagawara A,** Shimahara Y, Yamaoka Y, Fukumoto M. Comprehensive allelotyping of human intrahepatic cholangiocarcinoma. *Clin Cancer Res* 7:2648-2655, 2001.
177. Miwa M, Kanai M, Binlian S, Hirai M, Xin X, Saito S, Hanai S, Kaneko M, **Nakagawara A,** Uchida K. DNA methylation in neuroblastoma. In “*DNA methylation and cancer*”, Eds. Hirohashi S, Jones PA, Miwa M, Saya H, Ushijima T., pp104-106, 2001. (Extended Abstract for the 31st International Symposium of the Princess Takamatsu Cancer Research Fund, Tokyo, 2000)
178. Shinagawa T, Nomura T, Colmenares C, Ohira M, **Nakagawara A,** Ishii S. Increased susceptibility to tumorigenesis of *ski*-deficient heterozygous mice. *Oncogene* 20: 8100-8108, 2001.
179. **Nakagawara A.** TRK. In *Encyclopedic Reference of Cancer*, Ed. by Schwab M. 2001. Springer, Heidelberg, pp922-923.
180. Yoshida I, Koide S, **Nakagawara A,** Tsuji A, Matsuda Y. Proprotein convertase PACE4 is a target gene of the basic helix-loop-helix transcription factor hASH-1. *Biochem J* 360: 683-689, 2001.
181. Ogita J, Isogai E, Sudo H, Sakiyama S, **Nakagawara A,** Koseki H. Expression of the *Dan* gene during chicken embryonic development. *Mech. Dev.* 109: 363-365, 2001.
182. Kitanaka C, Kato K, Ijiri R, Sakurada K, Tomiyama A, Noguchi K, Nagashima Y, **Nakagawara A,** Momoi T, Yokoyama S, Toyoda Y, Kigasawa H, Nishi T, Tanaka Y, Kuchino Y. Increased Ras expression and caspase-independent neuroblastoma cell death: Possible mechanism of spontaneous neuroblastoma regression. *J. Natl. Cancer Inst.* 94: 358-368, 2002.
183. Ohtori S, Yamamoto T, Ino H, Hanaoka E, Shinbo J, Ozaki T, Takada N, Nakamura Y, Chiba T, **Nakagawara A,** Sakiyama S, Sakashita Y, Takahashi K, Tanaka K, Yamagata M, Yamazaki M, Shimizu S, Moriya H. Differential screening-selected gene aberrative in neuroblastoma protein modulates inflammatory pain in the spinal dorsal horn. *Neuroscience* 110: 579-586, 2002.
184. Ichimiya S, Kojima T, Momota H, Kondo N, Ozaki T, **Nakagawara A,** Toribio ML, Imamura M, Sato N. p73 is expressed in human thymic epithelial cells. *J. Histochem. Cytochem.* 50: 455-462, 2002.

185. Nakagawa T, Takahashi M, Ozaki T, Watanabe K, Todo S, Mizuguchi H, Hayakawa T, **Nakagawara A**. Autoinhibitory regulation of p73 by $\Delta Np73$ to modulate cell survival and death through p73-specific target element within the $\Delta Np73$ promoter. *Mol. Cell. Biol.* 22: 2575-2585, 2002.
186. Takahashi M, Seki N, Ozaki T, Kato M, Kuno T, Nakagawa T, Watanabe K, Miyazaki K, Ohira M, Hayashi S, Hosoda M, Tokita H, Mizuguchi H, Hayakawa T, Todo S, **Nakagawara A**. Identification of the p33^{ING1}-regulated genes which include *cyclin B1* and proto-oncogene *DEK* by using cDNA microarray in a mouse mammary epithelial cell line NMuMG. *Cancer Res.* 62: 2203-2209, 2002.
187. Stros M, Ozaki T, Bacikova A, Kageyama H, **Nakagawara A**. HNGB1 and HMGB2 cell-specifically down-regulate the p53- and p73-dependent sequence-specific transactivation from the human *Bax* gene promoter. *J. Biol. Chem.* 277: 7157-7164, 2002.
188. Watanabe K, Ozaki T, Nakagawa T, Miyazaki K, Takahashi M, Hosoda M, Hayashi S, Todo S, **Nakagawara A**. Physical interaction of p73 with c-myc and MM1, a c-myc-binding protein, and modulation of the p73 function. *J. Biol. Chem.* 277: 15113-15123, 2002.
189. Shinbo J, Ozaki T, Nakagawa T, Watanabe K, Nakamura Y, Yamazaki M, Moriya H, **Nakagawara A**, Sakiyama S. p73-dependent expression of DAN during cisplatin-induced cell death and osteoblast differentiation. *Biochem. Biophys. Res. Commun.* 295: 501-507, 2002.
190. Takayasu H, Motoi T, Kanamori Y, Kitano Y, Nakanishi H, Tange T, **Nakagawara A**, Hashizume K. Two case reports of childhood liver cell adenomas harboring β -catenin abnormalities. *Hum. Pathol.* 33: 852-855, 2002.
191. Iuchi T, Namba H, Iwadate Y, Shishikura T, Kageyama H, Nakamura Y, Ohira M, Tamaura A, Osato K, Sakiyama S, **Nakagawara A**. Identification of the small interstitial deletion at chromosome 1p34-p35 and its association with poor outcome in oligodendroglial tumors. *Genes Chromosomes Cancer* 35: 170-175, 2002.
192. Tomizawa M, Wang Y-Q, Ebara M, Saisho H, Watanabe K, **Nakagawara A**, Tagawa M. Decreased expression of the CCAAT/enhancer binding protein alpha gene involved in hepatocyte proliferation in human hepatocellular carcinomas. *Int. J. Mol. Med.* 9: 597-600, 2002.
193. Muramatsu T, Sakuma S, Ikematsu S, Shimada H, Okamoto K, Muramatsu H, Kadomatsu K, **Nakagawara A**. Midkine, a heparin-binding growth factor, is *overexpressed in human carcinomas*, and is useful as a tumor marker. In "Basic and clinical research on tumor markers", Eds. J. Carl Barrett, Kohzoh Imai, Tadao Kakizoe, John E. Shively, Ken Yamaguchi, pp42-47, 2002. (Extended Abstract for the 32nd International Symposium of the Princess Takamatsu Cancer Research Fund, Tokyo, 2001)
194. Tomioka N, Kobayashi H, Kageyama H, Ohira M, Nakamura Y, Sasaki F, Todo S, **Nakagawara A**, Kaneko Y. Chromosomes that show partial loss or gain in near-diploid tumors coincide with chromosomes that show whole loss or gain in near-triploid tumors: Evidence suggesting the involvement of the same genes in the tumorigenesis of high- and low-risk neuroblastomas. *Genes Chromosomes Cancer* 36:139-150, 2003.
195. Tomizawa M, Watanabe K, Saisho H, **Nakagawara A**, Tagawa M. Down-regulated expression of the CCAAT/enhancer binding protein a and b genes in human hepatocellular carcinoma: a possible prognostic marker. *Anticancer Res.* 23: 351-354, 2003.
196. Kawamoto T, Ohira M, Hamano S, Hori T, **Nakagawara A**. High expression of the novel endothelin-converting enzyme genes, *Nbla03145/ECEL1 α* and β , is associated with favorable prognosis in human neuroblastomas. *Int. J. Oncol.* 22:815-822, 2003.

197. Saito-Ohara F, Imoto I, Inoue J, **Nakagawara A**, Sugimoto T, Inazawa J. PPM1D is a potential target for 17q gain in neuroblastoma. *Cancer Res.* 63: 1876-1883, 2003.
198. Sakiyama S, Yu L, Tomizawa M, Shimada H, Kadomatsu K, Muramatsu T, Ikematsu S, **Nakagawara A**, Tagawa M. Utilization of the promoter region of the midkine gene as a tool to drive therapeutic genes in a tumor specific manner. *Advan. Enzyme Regul.* 43:57-66, 2003.
199. Carninci P, Waki K, Shiraki T, Konno H, Shibata K, Itoh M, Aizawa K, Arakawa T, Ishii Y, Sasaki D, Bono H, Kondo S, Sugahara Y, Saito R, Osato N, Fukuda S, Sato K, Watahiki A, Hirozane-Kishikawa T, Nakamura M, Shibata Y, Yasunishi A, Kikuchi N, Yoshiki A, Kusakabe M, Gustincich S, Beisel K, Pavan W, Aidinis V, **Nakagawara A**, Held W. A., Iwata H, Kono T, Nakauchi H, Lyons P, Wells C, Hume D. A., Fagiolini M, Hensch T. K., Brinkmeier M, Camper S, Muramatsu M, Okazaki Y, Kawai J, Hayashizaki Y. Targeting a complex transcriptome: The construction of the mouse full-length cDNA encyclopedia. *Genome Res.* 13:1273-1289, 2003.
200. Ozaki T, Watanabe K, Nakagawa T, Miyazaki K, Takahashi M, **Nakagawara A**. Function of p73, not of p53, is inhibited by the physical interaction with RACK1 and its inhibitory effect is counteracted by pRB. *Oncogene* 22:3231-3242, 2003.
201. Okamoto Y, Ozaki T, Miyazaki K, Aoyama M, Miyazaki M, **Nakagawara A**. UbcH10 is the cancer-related E2 ubiquitin conjugating enzyme. *Cancer Res.* 63:4167-4173, 2003.
202. Ikematsu S, **Nakagawara A**, Nakamura Y, Sakuma S, Wakai K, Muramatsu T, Kadomatsu K. Correlation of elevated level of serum midkine with poor prognostic factors of human neuroblastomas. *Br. J. Cancer* 88:1522-1526, 2003.
203. Nakagawa T, Takahashi M, Ozaki T, Watanabe K, Hayashi S, Hosoda M, Todo S, **Nakagawara A**. Negative autoregulation of p73 and p53 by $\Delta Np73$ in regulating differentiation and survival of human neuroblastoma cells. *Cancer Lett.* 197:105-109, 2003.
204. Ohira M, Morohashi A, Nakamura Y, Isogai E, Furuya K, Hamano S, Machida T, Aoyama M, Fukumura M, Miyazaki K, Suzuki Y, Sugano S, Hirato J, **Nakagawara A**. Neuroblastoma Oligo-capping cDNA Project : Toward the Understanding of the Genesis and Biology of Neuroblastoma *Cancer Lett.* 197:63-68, 2003.
205. Ohira M, Morohashi A, Inuzuka H, Shishikura T, Kawamoto T, Kageyama H, Nakamura Y, Isogai E, Takayasu H, Sakiyama S, Suzuki Y, Sugano S, Goto T, Sato S, **Nakagawara A**. Expression profiling and characterization of 4,200 genes cloned from primary neuroblastomas: Identification of 305 genes differentially expressed between favorable and unfavorable subsets. *Oncogene* 22:5525-5536, 2003.
206. Miyazaki K, Ozaki T, Kato C, Hanamoto T, Fujita T, Irino S, Watanabe K, Nakagawa T, **Nakagawara A**. A novel HECT-type E3 ubiquitin ligase, NEDL2, stabilizes p73 and enhances its transcriptional activity. *Biochem. Biophys. Res. Commun.* 308:106-113, 2003.
207. Nakamura Y, Ozaki T, Koseki H, **Nakagawara A**, Sakiyama S. Accumulation of p27^{KIP1} is associated with BMP2-mediated growth arrest and neuronal differentiation of human neuroblastoma-derived cell lines. *Biochem. Biophys. Res. Commun.* 307:206-213, 2003.
208. **Nakagawara A**. Neural crest development and neuroblastoma: the genetic and biological link. In NGF and Related Molecules in Health and Disease, Ed. By Luigi Aloe and Laura Calza, Progress in Brain Research Vol. 146, 2004, *Elsevier Science Publisher*, pp233-242.
209. **Nakagawara A**, Ohira M. Comprehensive genomics linking between neural development and cancer: Neuroblastoma as a model. In Special Issue: Neural development and cancer. *Cancer Lett.* 204:23-224, 2004.

210. Miyazaki K, Fujita T, Ozaki T, Kato C, Kurose Y, Sakamoto M, Kato S, Goto T, Itoyama Y, Aoki M, **Nakagawara A**. NEDL1, a novel ubiquitin-protein isopeptide ligase for Dishevelled-1, targets mutant superoxide dismutase-1. *J. Biol. Chem.* 279: 11327-11335, 2004.
211. Hamano S, Ohira M, Isogai E, Nakada K, **Nakagawara A**. Identification of novel human neuronal leucine-rich repeat (hNLRR) family genes and inverse association of expression of *Nbla10449/hNLRR-1* and *Nbla10677/hNLRR-3* with the prognosis of primary neuroblastomas. *Int. J. Oncol.* 24:1457-1466, 2004
212. Ohtori S, Isogai E, Hasue F, Ozaki T, Nakamura Y, **Nakagawara A**, Koseki H, Yuasa S, Hanaoka E, Shinbo J, Yamamoto T, Chiba H, Yamazaki M, Moriya H, Sakiyama S. Reduced inflammatory pain in mice deficient in the differential screening-selected gene abrrative in neuroblastoma. *Mol. Cell. Neurosci.* 25:504-514, 2004.
213. Wang YQ, Seimiya M, Kawamura K, Yu L, Ogi T, Takenaga K, Shishikura T, **Nakagawara A**, Sakiyama S, Tagawa M and O-Wang J. Elevated expression of DNA polymerase κ in human lung cancer is associated with p53 inactivation: negative regulation of POLK promoter activity by p53. *Int. J. Oncol.* 25:161-165, 2004.
214. Ando K, Ozaki T, Yamamoto H, Furuya K, Hosoda M, Hayashi S, Fukuzawa M, **Nakagawara A**. Polo-like kinase 1 (Plk1) inhibits p53 function by physical interaction and phosphorylation. *J. Biol. Chem.* 279:25549-25561. 2004.
215. Hiyama E, Yamaoka H, Matsunaga T, Hayashi Y, Ando H, Suita S, Horie H, Kaneko M, Sasaki F, Hashizume K, **Nakagawara A**, Ohnuma N, Yokoyama T. High expression of telomerase is an independent prognostic indicator of poor outcome in hepatoblastoma. *Br. J. Cancer* 91:972-979, 2004.
216. Yamada S, Ohira M, Horie H, Ando K, Takayasu H, Suzuki Y, Sugano S, Matsunaga T, Hiyama E, Hayashi Y, Watanabe Y, Suita S, Kaneko M, Sasaki F, Hashizume K, Ohnuma N, **Nakagawara A**. Expression profiling and differential screening between hepatoblastomas and the corresponding normal livers: Identification of high expression of the *Plk1* oncogene as a poor-prognostic indicator of hepatoblastomas. *Oncogene* 23:5901-5911, 2004.
217. Takahashi M, Ozaki T, Todo S, **Nakagawara A**. Decreased expression of the candidate tumor suppressor gene ING1 is associated with poor prognosis in advanced neuroblastomas. *Oncol Rep.* 12:811-816, 2004.
218. Kato C, Miyazaki K, Nakagawa A, Ohira M, Nakamura Y, Ozaki T, Imai T, **Nakagawara A**. Low expression of human tubulin tyrosine ligase and suppressed tubulin tyrosination/detyrosination cycle are associated with impaired neuronal differentiation in neuroblastomas with poor prognosis. *Int. J. Cancer* 112:365-375, 2004.
219. **Nakagawara A**. Chapter 5. Molecular and developmental biology of neuroblastoma. In Neuroblastoma, Eds. N-K. Cheung & S. Cohn, 2005, *Springer-Verlag, Heidelberg*, pp41-53.
220. Kramer S, Ozaki T, Miyazaki K, Kato C, Hanamoto T, **Nakagawara A**. Protein stability and function of p73 are modulated by a physical interaction with RanBPM in mammalian cultured cells. *Oncogene* 24:938-944, 2005
221. Abe M, Ohira M, Kaneda A, Yagi Y, Yamamoto S, Kitano Y, Takato T, **Nakagawara A**, Ushijima T. CpG island methylator phenotype is a strong determinant of poor prognosis in neuroblastomas. *Cancer Res.* 65:828-834, 2005
222. Ohira M, Oba S, Nakamura Y, Hirata T, Ishii S, **Nakagawara A**. A review of DNA microarray analysis of human neuroblastomas. *Cancer Lett.* 228:5-11, 2005

223. Lin L, Ozaki T, Takada Y, Kageyama H, Nakamura Y, Hata A, Zhang J-H, Simonds W, **Nakagawara A**, Koseki H. Topors, a p53 and topoisomerase I-binding RING finger protein, is a co-activator of p53 in growth suppression induced by DNA damage. *Oncogene* 24:3385-3396, 2005
224. Ozaki T, Hosoda M, Miyazaki K, Hayashi S, Watanabe K, Nakagawa T, **Nakagawara A**. Functional implication of p73 protein stability in neuronal cell survival and death. *Cancer Lett.* 228:29-35, 2005
225. Ohira M, Oba S, Nakamura Y, Isogai E, Kaneko S, Nakagawa A, Hirata T, Kubo H, Goto T, Yamada S, Yoshida Y, Fuchioka M, Ishii S, **Nakagawara A**. Expression profiling using a tumor-specific cDNA microarray predicts the prognosis of intermediate-risk neuroblastomas. *Cancer Cell* 7:337-350, 2005
226. Hanamoto T, Ozaki T, Furuya K, Hosoda M, Hayashi S, Nakanishi M, Yamamoto H, Kikuchi H, Todo S, **Nakagawara A**. Identification of protein kinase A catalytic subunit beta as a novel binding partner of p73 and regulation of p73 function. *J. Biol. Chem.* 280:16665-16675, 2005
227. Gotoh T, Hosoi H, Iehara T, Kuwahara Y, Osone S, Tsuchiya K, Kuroda H, Ohira M, **Nakagawara A**, Sugimoto T. Prediction of MYCN amplification in neuroblastoma using serum DNA and real-time quantitative PCR. *J. Clin. Oncol.* 23: 5205-5210, 2005
228. Osajima-Hakomori Y, Miyake I, Ohira M, **Nakagawara A**, Nakagawa A, Sakai R. Biological role of anaplastic lymphoma kinase in neuroblastoma. *Am J Pathol.* 167:213-222, 2005
229. Aoyama M, Ozaki T, Inuzuka H, Tomotsune D, Hirato J, Okamoto Y, Tokita H, Ohira M, **Nakagawara A**. LMO3 interacts with neuronal transcription factor, HEN2, and acts as an oncogene in neuroblastoma. *Cancer Res.* 65:4587-4597, 2005
230. Chen H, Suzuki M, Nakamura Y, Ohira M, Ando S, Iida T, Nakajima T, **Nakagawara A**, Kimura H. Aberrant methylation of *FBN2* correlates with progression of human non-small cell lung cancer. *Lung Cancer* 50:43-49, 2005
231. **Nakagawara A**, Sawada T, Brodeur GM, Matthay KK. Professor Yoshiaki Tsuchida, MD, PhD (1936-2005). *Pediatr. Blood Cancer.* 45:609-610, 2005
232. Okabe-Kado J, Kasukabe T, Honma Y, Hanada R, **Nakagawara A**, Kaneko Y. Clinical significance of serum NM23-H1 protein in neuroblastoma. *Cancer Sci.* 96:653-660, 2005
233. Ozaki T, **Nakagawara A**. p73, a sophisticated p53 family member in the cancer world. *Cancer Sci.* 96:729-737, 2005
234. Hosoda M, Ozaki T, Miyazaki K, Hayashi S, Furuya K, Watanabe K, Nakagawa T, Hanamoto T, Todo S, **Nakagawara A**. UFD2a mediates the proteasomal turnover of p73 without promoting p73 ubiquitination. *Oncogene* 24:7156-7169, 2005
235. Koshikawa N, Maejima C, Miyazaki K, **Nakagawara A**, Takenaga K. Hypoxia selects for high-metastatic Lewis lung carcinoma cells overexpressing Mcl-1 and exhibiting reduced apoptotic potential in solid tumors. *Oncogene* 25:917-928, 2006
236. Kaneko Y, Kobayashi H, Watanabe N, Tomioka N, **Nakagawara A**. Biology of neuroblastomas that were found by mass screening at 6 months of age in Japan. *Pediatr. Blood Cancer* 46:285-291, 2006
237. Inamori K, Gu J, Ohira M, Kawasaki A, Nakamura Y, Nakagawa T, Kondo A, Miyoshi E, **Nakagawara A**, Taniguchi N. High expression of N-acetylglucosaminyltransferase V in favorable neuroblastomas: Involvement of its effect on apoptosis. *FEBS Lett.* 580:627-632, 2006

238. Isono K, Nemoto K, Li Y, Takada Y, Suzuki R, Katsuki M, **Nakagawara A**, Koseki H. Overlapping roles for homeodomain-interacting protein kinases Hipk1 and Hipk2 in the mediation of cell growth in response to morphogenetic and genotoxic signals. *Mol. Cell. Biol.* 26:2758-2771, 2006
239. Machida T, Fujita T, Ooo M L, Ohira M, Isogai E, Mihara M, Hirato J, Tomotsune D, Hirata T, Fujimori M, Adachi W, **Nakagawara A**. Decreased expression of pro-apoptotic *BMCC1*, a novel gene with the *BNIP2* and *Cdc42GAP* homology (BCH) domain, is associated with poor prognosis in human neuroblastomas. *Oncogene* 25:1931-1942, 2006
240. Niizuma H, Nakamura Y, Ozaki T, Ohira M, Isogai E, Kageyama H, Imaizumi M, **Nakagawara A**. Bcl-2 is a key regulator for the retinoic acid-induced apoptotic cell death in neuroblastoma. *Oncogene* 25:5046-5055, 2006.
241. Chen H, Suzuki M, Nakamura Y, Ohira M, Ando S, Iida T., Nakajima T, **Nakagawara A**, Kimura H. Aberrant methylation of *RASGRF2* and *RASSF1A* in human non-small cell lung cancer. *Oncol. Rep.* 15:1281-1285, 2006
242. Sang M, Li Y, Ozaki T, Ono S, Ando K, Yamamoto H, Koda T, Geng C, **Nakagawara A**. p73-dependent induction of 14-3-3 σ increases the chemo-sensitivity of drug-resistant human breast cancers. *Biochem. Biophys. Res. Commun.* 347:327-333, 2006
243. Hayashi S, Ozaki T, Yoshida K, Hosoda M, Todo S, Akiyama S, **Nakagawara A**. p73 and MDM2 confer the resistance of epidermoid carcinoma to cisplatin by blocking p53. *Biochem. Biophys. Res. Commun.* 347:60-66, 2006
244. Ozaki T, Li Y, Kikuchi H, Tomita T, Iwatsubo T, **Nakagawara A**. The intracellular domain of the amyloid precursor protein (AICD) enhances the p53-mediated apoptosis. *Biochem. Biophys. Res. Commun.* 351:57-63, 2006
245. Kikuchi H, Ozaki T, Furuya K, Hanamoto T, Nakanishi M, Yamamoto H, Yoshida K, Todo S, **Nakagawara A**. NF-kB regulates the stability and activity of p73 by inducing its proteolytic degradation through a ubiquitin-dependent proteasome pathway. *Oncogene* 25:7608-7617, 2006
246. Okahara F, Itoh K, **Nakagawara A**, Murakami M, Kanaho Y, Maehara T. Critical role of PICT1, a tumor suppressor candidate, in phosphatidylinositol 3,4,5-triphosphate signals and tumorigenic transformation. *Mol. Biol. Cell* 17:4888-4895, 2006
247. Abe M, Westermann F, **Nakagawara A**, Takato T, Schwab M, Ushijima T. Marked and independent prognostic significance of the CpG island methylator phenotype in neuroblastomas. *Cancer Lett.* 247:253-258, 2007
248. Tomizawa M, Horie H, Yamamoto H, Matsunaga T, Sasaki F, Hashizume K, Hiyama E, Kaneko M, Suita S, Ando H, Hayashi Y, Ohnuma N, **Nakagawara A**. Reciprocal expression of CCAAT/enhancer binding proteins α and β in hepatoblastomas and its prognostic significance. *Oncol. Rep.* 17:341-344, 2007
249. Kaneko S, Ohira M, Nakamura Y, Isogai E, **Nakagawara A**, Kaneko M. Relationship of DDX1 and NAG gene amplification/overexpression to the prognosis of patients with MYCN-amplified neuroblastoma. *J. Cancer Res. Clin. Oncol.* 133:185-192, 2007
250. Antonelli A, Lenzi L, **Nakagawara A**, Ozaki T, Chiaretti A, Aloe L. Tumor suppressor proteins are differentially affected in human ependyoblastoma and medulloblastoma cells exposed to nerve growth factor. *Cancer Investigation* 25:94-101, 2007
251. Yamamoto H, Ozaki T, Nakanishi M, Kikuchi H, Yoshida K, Horie H, Kuwano H, **Nakagawara A**. Oxidative stress induces p53-dependent apoptosis in hepatoblastoma cell through its nuclear translocation. *Genes to Cells* 12:461-471, 2007

252. Nakamura Y, Ozaki T, Niizuma H, Ohira M, Kamijo T, **Nakagawara A**. Functional characterization of a new p53 mutant generated by homozygous deletion in a neuroblastoma cell line. *Biochem. Biophys. Res. Commun.* 354:892-898, 2007
253. Nakanishi H, Ozaki T, Nakamura Y, Hashizume K, Iwanaka T, **Nakagawara A**. Purification of human primary neuroblastomas by magnetic beads and their in vitro culture. *Oncol. Rep.* 17:1315-1320, 2007
254. Nimura Y, Kawata T, Uzawa K, Okamura J, Liu C, Saito M, Shimada H, Seki N, **Nakagawara A**, Ito H, Ochiai T, Tanzawa H. Silencing Ku80 using small interfering RNA enhanced radiation sensitivity in vitro and in vivo. *Int. J. Oncol.* 30:1477-1484, 2007
255. Takahashi M, Ozaki T, Takahashi A, Miyauchi M, Ono S, Takada N, Koda T, Todo S, Kamijo T, **Nakagawara A**. DFF45/ICAD restores cisplatin-induced nuclear fragmentation but not DNA cleavage in DFF45-deficient neuroblastoma cells. *Oncogene* 26:5669-5673, 2007
256. Furuya K, Ozaki T, Hanamoto T, Hosoda M, Hayashi S, Barker PA, Takano K, Matsumoto M, **Nakagawara A**. Stabilization of p73 by nuclear I κ B kinase- α mediates cisplatin-induced apoptosis. *J. Biol. Chem.* 282:18365-18378, 2007
257. Nakanishi M, Ozaki T, Yamamoto H, Hanamoto T, Kikuchi H, Furuya K, Asaka M, Delia D, **Nakagawara A**. NFBD1/MDC1 associates with p53 and regulates its function at the crossroad between cell survival and death in response to DNA damage. *J. Biol. Chem.* 282:22993-23004, 2007
258. Iwao-Koizumi K, Maekawa K, Nakamura Y, Saito S, Kawamoto S, **Nakagawara A**, Kato K. A novel technique for measuring variations in DNA copy-number: competitive genomic polymerase chain reaction. *BMC Genomics* 8:206.2007
259. Ichikawa T, Suenaga Y, Koda T, Ozaki T, **Nakagawara A**. TAp63-dependent induction of growth differentiation factor 15 (GDF15) plays a critical role in the regulation of keratinocyte differentiation. *Oncogene* 27:409-420, 2008
260. Tomioka N, Oba S, Ohira M, Misra A, Fridlyand J, Ishii S, Nakamura Y, Isogai E, Hirata T, Yoshida Y, Todo S, Kaneko Y, Albertson DG, Pinkel D, Feuerstein BG, **Nakagawara A**. Novel risk stratification of patients with neuroblastoma by genomic signature, which is independent of molecular signature. *Oncogene* 27:441-449, 2008
261. Yoshida K, Ozaki T, Furuya K, Nakanishi M, Kikuchi H, Yamamoto H, Ono S, Koda T, Omura K, **Nakagawara A**. ATM-dependent nuclear accumulation of IKK- α plays an important role in the regulation of p73-mediated apoptosis in response to cisplatin. *Oncogene* 27:1183-1188.2008
262. Arai H, Ozaki T, Niizuma H, Nakamura Y, Ohira M, Sugita K, **Nakagawara A**. ERAP140/Nbla10993 is a novel favorable prognostic indicator for neuroblastoma and induced in response to retinoic acid. *Oncol. Rep.* 19:1381-1388, 2008
263. Kurata K, Yanagisawa R, Ohira M, Kitagawa M, **Nakagawara A**, Kamijo T. Stress via p53 pathway causes apoptosis by mitochondrial Noxa up-regulation in doxorubicin-treated neuroblastoma cells. *Oncogene* 27:741-754, 2008
264. Bu Y, Suenaga Y, Ono S, Koda T, Son F, **Nakagawara A**. Ozaki T, Sp1-mediated transcriptional regulation of NFBD1/MDC1 plays a critical role in DNA damage response pathway. *Genes Cells.* 13:53-66, 2008
265. Okoshi R, Ozaki T, Yamamoto H, Ando K, Koida N, Ono S, Koda T, Kamijo T, **Nakagawara A**, Kizaki H. Activation of AMP-activated protein kinase induces p53-dependent apoptotic cell death in response to energetic stress. *J. Biol. Chem.* 283:3979-3987, 2008

266. Li Y, Ozaki T, Kikuchi H, Yamamoto H, Ohira M, **Nakagawara A**. A novel HECT-type E3 ubiquitin protein ligase NEDL1 enhances the p53-mediated apoptotic cell death in its catalytic activity-independent manner. *Oncogene*. 27:3700-3709, 2008
267. Ichikawa T, Suenaga Y, Koda T, Ozaki T, **Nakagawara A**. Δ Np63/BMP-7-dependent expression of matrilin-2 is involved in keratinocyte migration in response to wounding. *Biochem. Biophys. Res. Commun.* 369:994-1000, 2008
268. Honda S, Haruta M, Sugawara W, Sasaki F, Ohira M, Matsunaga T, Yamaoka H, Horie H, Ohnuma N, **Nakagawara A**, Hiyama E, Todo S, Kaneko Y. The methylation status of RASSF1A promoter predicts responsiveness to chemotherapy and eventual cure in hepatoblastoma patients. *Int. J. Cancer* 123:1117-1125, 2008.
269. Wang H, Ozaki T, Hossain MS, Nakamura Y, Kamijo T, Xue X, **Nakagawara A**. A newly identified dependence receptor UNC5H4 is induced during DNA damage-mediated apoptosis and transcriptional target of tumor suppressor p53. *Biochem. Biophys. Res. Commun.* 370:594-8, 2008
270. Abe M, Watanabe N, McDonell N, Takato T, Ohira M, **Nakagawara A**, Ushijima T. Identification of genes targeted by CpG island methylator phenotype in neuroblastomas, and their possible integrative involvement in poor prognosis.. *Oncology* 74:50-60, 2008.
271. Koida N, Ozaki T, Yamamoto H, Ono S, Koda T, Ando K, Okoshi R, Kamijo T, Omura K, **Nakagawara A**. Inhibitory role of Plk1 in the regulation of p73-dependent apoptosis through physical interaction and phosphorylation. *J. Biol. Chem.* 283:8555-63, 2008
272. Inoue K-I, Ozaki T, Nakanishi M, Kikuchi H, Yamamoto H, Todo S, **Nakagawara A**. NFBD1/MDC1 stabilizes oncogenic MDM2 to contribute to cell fate determination in response to DNA damage. *Biochem Biophys Res Commun.* 371:829-833, 2008
273. Nakagawa H, Ohira M, Hayashi S, Abe S, Saito S, Nagahori N, Monde K, Shinohara Y, Fujitani N, Kondo H, Akiyama S, **Nakagawara A**, Nishimura S. Alterations in the glycoform of cisplatin-resistant human carcinoma cells is caused by defects in the endoplasmic reticulum-associated degradation system. *Cancer Lett.* 270:295-301, 2008
274. Hossain MS, Ozaki T, Wang H, Nakagawa A, Takenobu H, Ohira M, Kamijo T, **Nakagawara A**. N-MYC promotes cell proliferation through a direct transactivation of neuronal leucine-rich repeat protein-1 (NLRRL1) gene in neuroblastoma. *Oncogene* 27:6075-6082, 2008
275. Ando K, Ohira M, Ozaki T, Nakagawa A, Akazawa K, Suenaga Y, Nakamura Y, Koda T, Kamijo T, Murakami Y, **Nakagawara A**. Expression of TSLC1, a candidate tumor suppressor gene mapped to chromosome 11q23, is down-regulated in unfavorable neuroblastoma without promoter hypermethylation. *Int. J. Cancer* 123:2087-2094, 2008
276. Munirajan AK, Ando K, Mukai A, Takahashi M, Suenaga Y, Ohira M, Koda T, Hirota T, Ozaki T, **Nakagawara A**. KIF1B β functions as a haploinsufficient tumor suppressor gene mapped to chromosome 1p36.2 by inducing apoptotic cell death. *J. Biol. Chem.* 283:24426-24434, 2008
277. Chen Y, Takita J, Choi YL, Kato M, Ohira M, Sanada M, Soda M, Kikuchi A, Igarashi T, **Nakagawara A**, Hayashi Y, Mano H, Ogawa S. Novel oncogenic mutations of ALK kinase in neuroblastoma. *Nature* 455:971-974, 2008.
278. Ikematsu S, **Nakagawara A**, Nakamura Y, Ohira M, Shinjo M, Kishida S, Kadomatsu K. Plasma midkine level is a prognostic factor for human neuroblastoma. *Cancer Sci.* 99:2070-2074, 2008

279. Honda S, Arai Y, Haruta M, Sasaki F, Ohira M, Yamaoka H, Horie H, **Nakagawara A**, Hiyama E, Todo S, Kaneko Y. Loss of imprinting of IGF2 correlates with hypermethylation of the H19 differentially methylated region in hepatoblastoma. *Br. J. Cancer* 99:1891-1899, 2008
280. Fujita T, Ikeda H, Kawasaki K, Taira N, Ogasawara Y, **Nakagawara A**, Doihara H. Clinicopathological relevance of UbcH10 in breast cancer. *Cancer Sci.* 2008 (Epub ahead of print)
281. Miyake I, Ohira M, **Nakagawara A**, Sakai R. Distinct role of ShcC docking protein in the differentiation of neuroblastoma. *Oncogene* 28:662-673, 2009
282. Okoshi R, Ando K, Suenaga Y, Kizaki H, **Nakagawara A**, Ozaki T. Transcriptional regulation of tumor suppressor p53 by CREB/AMPK complex in response to glucose deprivation. *J. Biol. Chem.* (under revision)