

業 績 目 録

原 著 論 文

1. **Ozaki T**, Hoshikawa Y, Iida Y, Iwabuchi M.
Sequence analysis of the transcribed and 5' non-transcribed regions of the ribosomal RNA gene in *Dictyostelium discoideum*. **Nucleic Acids Res.**, 1984; **10**:4171-4184.
2. **Ozaki T**, Hasegawa M, Hamada Y, Tasaka M, Iwabuchi M, Takeuchi I. Molecular cloning of cell-type-specific cDNAs exhibiting new types of developmental regulation in *Dictyostelium discoideum*. **Cell Differ.**, 1988, **23**:119-124.
3. Tasaka M, Hasegawa M, **Ozaki T**, Iwabuchi M, Takeuchi I. Isolation and characterization of spore coat protein (sp96) gene of *Dictyostelium discoideum*. **Cell Differ. Development**, 1990, **31**:1-9.
4. Morio T, **Ozaki T**, Takeuchi I, Tasaka M.
Transcriptional regulation of cell-type-enriched genes during development in *Dictyostelium discoideum* analyzed by nuclear run-on assays. **Development Growth Differ.**, 1991, **33**:293-298.
5. Tasaka M, Hasegawa M, Nakata M, Orii H, **Ozaki T**, Takeuchi I. Protein binding and DNase-I- hypersensitive sites in the *cis*-acting regulatory region of the spore-coat SP96 gene of *Dictyostelium*. **Mec. Dev.** 1992, **36**:105-115.
6. **Ozaki T**, Sakiyama S. Lamin A gene expression is specifically suppressed in *v-src*-transformed cells. **FEBS Lett.**, 1992, **312**:165-168.
7. **Ozaki T**, Nakao H, Orii H, Morio T, Takeuchi I, Tasaka M. Developmental regulation of transcription of a novel prespore-specific gene (*Dp87*) in *Dictyostelium discoideum*. **Development**, 1993, **117**:1299-1308.
8. **Ozaki T**, Sakiyama S. Molecular cloning and characterization of a cDNA showing negative regulation in *v-src*-transformed 3Y1 rat fibroblasts. **Proc. Natl. Acad. Sci. USA**, 1993, **90**:2593-2597.
9. **Ozaki T**, Sakiyama S. Molecular cloning of rat calpactin I heavy-chain cDNA whose expression is induced in *v-src*-transformed rat culture cell lines. **Oncogene**, 1993, **8**: 1707-1710.
10. **Ozaki T**, Asai T, Sakiyama S. Molecular cloning of a unique complementary DNA of rat myosin regulatory light chain and its elevated expression in *v-src*-transformed rat culture cell lines. **Cell Growth Differ.**, 1993, **4**:939-945.
11. **Ozaki T**, Sakiyama S. Tumor-suppressive activity of *N03* gene product in *v-src*-transformed rat 3Y1 fibroblasts. **Cancer Res.**, 1994, **54**:646-648.
12. **Ozaki T**, Saijo M, Murakami K, Enomoto H, Taya Y, Sakiyama S. Complex formation between lamin A and the retinoblastoma gene product: Identification of the domain on lamin A required for its interaction. **Oncogene**, 1994, **9**: 2649-2653.
13. Enomoto H, **Ozaki T**, Takahashi E, Nomura N, Tabata S,

- Takahashi H, Ohnuma N, Tanabe M, Iwai J, Matsunaga T, Sakiyama S. Identification of human *DAN* gene, mapping to the putative neuroblastoma tumor suppressor locus. **Oncogene**, 1994, **9**:2785-2791.
14. Sakiyama S, **Ozaki T**, Enomoto H. Molecular cloning and characterization of a cDNA showing tumor-suppressive activity in v-*src*-transformed 3Y1 rat fibroblasts. **Advan Enzyme Regul.**, 1994, **34**:247-255.
 15. **Ozaki T**, Nakamura Y, Enomoto H, Hirose M, Sakiyama S. Overexpression of *DAN* gene product in normal rat fibroblasts causes retardation of the entry into the S phase. **Cancer Res.**, 1995, **55**:895-900.
 16. **Ozaki T**, Irie K, Sakiyama S. Molecular cloning and cell cycle-dependent expression of a novel gene that is homologous to *cdc37*. **DNA Cell Biol.**, 1995, **14**:1017-1023.
 17. **Ozaki T**, Sakiyama S. Sequence-specific DNA binding activity of rat *cdc37*-related gene product. **FEBS Lett.**, 1995, **375**:155-158.
 18. Kondo K, **Ozaki T**, Nakamura Y, Sakiyama S. *DAN* gene product has an affinity for Ni²⁺. **Biochem. Biophys. Res. Commun.**, 1995, **216**:209-215.
 19. **Ozaki T**, Ma J, Takenaga K, Sakiyama S. Cloning of mouse *DAN* cDNA and its down-regulation in transformed cells. **Jpn. J. Cancer Res.**, 1996, **87**:58-61.
 20. **Ozaki T**, Sakiyama S. Specific interaction of pRB with a rat genomic DNA fragment, REC11. **Biochem. Biophys. Res. Commun.**, 1996, **226**:237-241.
 21. **Ozaki T**, Sakiyama S. Interaction of rat *cdc37*-related protein with retinoblastoma gene product. **DNA Cell Biol.**, 1996, **15**:975-979.
 22. Matsuda Y, **Ozaki T**, Enomoto H, Saito T, Sakiyama S. Chromosome mapping of the mouse and rat homologues of the human *DAN* gene, *DIS1733E*. **Mammalian Genome**, 1996, **7**:709-710.
 23. **Ozaki T**, Sakiyama S. Identification of essential *cis*-acting regulatory elements for transcription of rat *DAN* gene. **DNA Cell Biol.** 1997; **16**:779-786.
 24. **Ozaki T**, Hishiki T, Toyama Y, Yuasa S, Nakagawara A, Sakiyama S. Identification of a new cellular protein which can interact specifically with DAN. **DNA Cell Biol.**, 1997, **16**:985-991.
 25. **Ozaki T**, Enomoto H, Nakamura Y, Kondo K, Seki N, Ohira M, Nomura N, Ohki M, Nakagawara A, Sakiyama S. The genomic analysis of human *DAN* gene. **DNA Cell Biol.**, 1997, **16**:1031-1039.
 26. **Ozaki T**, Kondo K, Nakamura Y, Ichimiya S, Nakagawara A, Sakiyama S. Interaction of DA41, a DAN-binding protein, with the epidermal growth factor-like protein, S(1-5). **Biochem. Biophys. Res. Commun.**, 1997, **237**:245-250.
 27. Nakamura Y, **Ozaki T**, Nakagawara A, Sakiyama S. A product of *DAN*, a novel candidate tumour suppressor gene, is secreted into culture medium and suppresses DNA synthesis. **Eur. J. Cancer.**, 1997, **33**:1986-1990.
 28. Nakamura Y, **Ozaki T**, Ichimiya S, Nakagawara A,

- Sakiyama S. Ectopic expression of DAN enhances the retinoic acid-induced neuronal differentiation in human neuroblastoma cell lines. **Biochem. Biophys. Res. Commun.**, 1998, **243**:722-726.
29. Hishiki T, Nimura Y, Isogai E, Kondo K, Ichimiya S, Nakamura Y, **Ozaki T**, Sakiyama S, Hirose M, Seki N, Takahashi H, Ohnuma N, Tanabe M, Nakagawara A. GDNF/NTN-induced differentiation and its enhanced by retinoic acid in primary human neuroblastomas expressing c-Ret, GFRA-1 and GFRA-2. **Cancer Res.**, 1998, **58**:2158-2165.
30. Uchimura K, Muramatsu H, Kaname T, Ogawa H, Yamakawa T, Fan Q-W, Mitsuoka C, Kannagi R, Habuchi O, Yokoyama I, Yamamura K, **Ozaki T**, Nakagawara A, Kadomatsu K, Muramatsu T. Human *N*-acetylglucosamine-6-*O*-sulfotransferase involved in the biosynthesis of β -sulfo sialyl lewis X: Molecular cloning, chromosomal mapping, and expression in various organs and tumor cells. **J. Biochem.**, 1998, **124**:670-678.
31. Hiwasa T, Nakamura Y, **Ozaki T**, Kondo K, Saido T, Nakagawara A, Sakiyama S. Down-regulation of protein kinase Ca and g and enhanced TPA-induced neurite formation in *DAN*-transfected neuroblastoma cells. **FEBS Lett.**, 1998, **440**:25-28.
32. Tomita S, **Ozaki T**, Taru H, Oguchi S, Takeda S, Yagi Y, Sakiyama S, Kirino Y, Suzuki T. Interaction of a neuron-specific protein containing PDZ domains with Alzheimer's amyloid precursor protein. **J. Biol. Chem.**, 1999, **274**:2243-2254.
33. Shishikura T, Ichimiya S, **Ozaki T**, Nimura Y, Kageyama H, Nakamura Y, Sakiyama S, Miyauchi M, Yamamoto N, Suzuki M, Nakajima N, Nakagawara A. Mutational analysis of the *p73* gene in human breast cancers. **Int. J. Cancer.**, 1999, **84**:321-325.
34. Takada N, **Ozaki T**, Ichimiya S, Todo S, Nakagawara A. Identification of a transactivation activity in the COOH-terminal region of p73 which is impaired in the naturally occurring mutants found in human neuroblastomas. **Cancer Res.**, 1999, **59**:2810-2814.
35. **Ozaki T**, Hanaoka E, Naka M, Nakagawara A, Sakiyama S. Cloning and characterization of rat *BAT3* cDNA. **DNA Cell Biol.**, 1999, **18**:503-512.
36. **Ozaki T**, Naka M, Takada N, Tada M, Sakiyama S, Nakagawara A. Deletion of the COOH-terminal region of p73a enhances both its transactivation function and DNA-binding activity but inhibits induction of apoptosis in mammalian cells. **Cancer Res.**, 1999, **59**:5902-5907.
37. Hanaoka E, **Ozaki T**, Ohira M, Nakamura Y, Suzuki M, Takahashi E, Moriya H, Nakagawara A, Sakiyama S. Molecular cloning and expression analysis of the human *DA41* gene and its mapping to chromosome 9q21.2-q21.3. **J. Hum. Genet.**, 2000, **45**:188-191.
38. Nagai M, Ichimiya S, **Ozaki T**, Seki N, Mihara M, Furuta S, Ohira M, Tomioka M, Nomura N, Sakiyama S,

- Kubo O, Takakura K, Hori T, Nakagawara A. Identification of the full-length *KIAA0591* gene encoding a novel kinesin-related protein which is mapped to the neuroblastoma suppressor gene locus at 1p36.2. **Int. J. Oncol.**, 2000, **16**: 907-916.
39. **Ozaki T**, Nagase T, Ichimiya S, Seki N, Ohira M, Nomura N, Takada N, Sakiyama S, Weber BL, Nakagawara A. NFBD1/KIAA0170 is a novel nuclear transcriptional transactivator with BRCT domain. **DNA Cell Biol.**, 2000, **19**:475-485.
40. **Ozaki T**, Nakamura Y, Hanaoka E, Nakagawara A, Sakiyama S. Overexpression of *DA41* in v-Ha-*ras*-3Y1 cells causes growth suppression. **Jpn. J. Cancer Res.**, 2000, **91**:987-993.
41. Hanaoka E, **Ozaki T**, Nakamura Y, Moriya H, Nakagawara A, Sakiyama S. Overexpression of DAN causes a growth suppression in p53-deficient SAOS-2 cells. **Biochem. Biophys. Res. Commun.**, 2000, **278**:20-26.
42. Ichimiya S, Nimura Y, Seki N, **Ozaki T**, Nagase T, Nakagawara A. Down-regulation of hASH1 is associated with the retinoic acid-induced differentiation of human neuroblastoma cell lines. **Med. Pediatr. Oncol.**, 2001, **36**:132-134.
43. Takayasu H, Horie H, Hiyama E, Matsunaga T, Hayashi Y, Watanabe Y, Suita S, Kaneko M, Sasaki F, Hashizume K, **Ozaki T**, Furuichi K, Tada M, Ohnuma N, Nakagawara A. Frequent deletions and mutations of the *b-catenin* gene are associated with overexpression of *cyclin D1* and *fibronectin* and poorly-differentiated histology in childhood hepatoblastoma. **Clin. Cancer Res.**, 2001, **7**:901-908.
44. Naka M, **Ozaki T**, Takada N, Takahashi M, Shishikura T, Sakiyama S, Tada M, Todo S, Nakagawara A. Functional characterization of naturally occurring mutants (P405R and P425L) of p73a and p73b found in neuroblastoma and lung cancer. **Oncogene**, 2001, **20**:3568-3572.
45. Stros M, **Ozaki T**, Ikova AB, Kageyama H, Nakagawara A. HMGB1 and HMGB2 cell-specifically down-regulate the p53- and p73-dependent sequence-specific transactivation from the human *Bax* gene promoter. **J. Biol. Chem.**, 2002, **277**:7157-7164.
46. Watanabe K, **Ozaki T**, Nakagawa T, Miyazaki K, Takahashi M, Hosoda M, Hayashi S, Todo S, Nakagawara A. Physical interaction of p73 with c-myc and MML, a c-myc-binding protein, and modulation of the p73 function. **J. Biol. Chem.**, 2002, **277**:15113-15123.
47. Takahashi M, Seki N, **Ozaki T**, Kato M, Kuno T, Nakagawa T, Watanabe K, Miyazaki K, Ohira M, Hayashi S, Hosoda M, Tokita H, Mizuguchi H, Hayakawa T, Todo S, Nakagawara A. Identification of the p33ING1-regulated genes which include *cyclin B1* and proto-oncogene *DEK* by using cDNA microarray in a mouse mammary epithelial cell line NMuMG. **Cancer Res.**, 2002, **62**:2203-2209.

48. Nakagawa T, Takahashi M, **Ozaki T**, Watanabe K, Todo S, Mizuguchi H, Hayakawa T, Nakagawara A. Autoinhibitory regulation of p73 by delta Np73 to modulate cell survival and death through p73-specific target element within the delta *Np73* promoter. **Mol. Cell. Biol.**, 2002, **22**:2575-2585.
49. Ichimiya S, Kojima T, Momota H, Kondo N, **Ozaki T**, Nakagawara A, Toribio ML, Imamura M, Sato N. p73 is expressed in human thymic epithelial cells. **J. Histochem. Cytochem.**, 2002, **50**:455-462.
50. Ohtori S, Yamamoto T, Ino H, Hanaoka E, Shinbo J, **Ozaki T**, Takada N, Nakamura Y, Chiba T, Nakagawara A, Sakiyama S, Sakashita Y, Takahashi K, Tanaka K, Yamagata M, Yamazaki M, Shimizu S, Moriya H. Differential screening-selected gene aberrative in neuroblastoma protein modulates inflammatory pain in the spinal dorsal horn. **Neuroscience**, 2002, **110**: 579-586.
51. Shinbo J, **Ozaki T**, Nakagawa T, Watanabe K, Nakamura Y, Yamazaki M, Moriya H, Nakagawara A, Sakiyama S. p73-dependent expression of DAN during cisplatin-induced cell death and osteoblast differentiation. **Biochem. Biophys. Res. Commun.**, 2002, **295**:501-507.
52. **Ozaki T**, Watanabe K, Nakagawa T, Miyazaki K, Takahashi M, Nakagawara A. Function of p73, not of p53, is inhibited by the physical interaction with RACK1 and its inhibitory effect is counteracted by pRB. **Oncogene**, 2003, **22**:3231-3242.
53. Nakamura Y, **Ozaki T**, Koseki H, Nakagawara A, Sakiyama S. Accumulation of p27KIP1 is associated with BMP2-induced growth arrest and neuronal differentiation of human neuroblastoma-derived cell lines. **Biochem. Biophys. Res. Commun.**, 2003, **307**:206-213.
54. Okamoto Y, **Ozaki T**, Miyazaki K, Aoyama M, Miyazaki M, Nakagawara A. UbcH10 is the cancer-related E2 ubiquitin conjugating enzyme. **Cancer Res.**, 2003, **63**: 4167-4173.
55. Miyazaki K, **Ozaki T**, Kato C, Hanamoto T, Fujita T, Irino S, Watanabe K, Nakagawa T, Nakagawara A. A novel HECT-type E3 ubiquitin ligase, NEDL2, stabilizes p73 and enhances its transcriptional activity. **Biochem. Biophys. Res. Commun.**, 2003, **308**:106-113.
56. Nakagawa T, Takahashi M, **Ozaki T**, Watanabe K, Hayashi S, Hosoda M, Todo S, Nakagawara A. Negative autoregulation of p73 and p53 by DeltaNp73 in regulating differentiation and survival of human neuroblastoma cells. **Cancer Lett.**, 2003, **197**: 105-109.
57. Miyazaki K, Fujita T, **Ozaki T**, Kato C, Kurose Y, Sakamoto M, Kato S, Goto T, Itoyama Y, Aoki M, Nakagawara A. NEDL1, a novel ubiquitin-protein isopeptide ligase for Dishevelled-1, targets mutant superoxide dismutase-1. **J. Biol. Chem.**, 2004, **279**: 11327-11335.
58. Ohtori S, Isogai E, Hasue F, **Ozaki T**, Nakamura Y,

- Nakagawara A, Koseki H, Yuasa S, Hanaoka E, Shinbo J, Yamamoto T, Chiba H, Yamazaki M, Moriya H, Sakiyama S. Reduced inflammatory pain in mice deficient in the differential screening-selected gene aberrative in neuroblastoma. **Mol. Cell Neurosci.**, 2004, **25**: 504-514.
59. Ando K, **Ozaki T**, Yamamoto H, Furuya K, Hosoda M, Hayashi S, Fukuzawa M, Nakagawara A. Polo-like kinase 1 (Plk1) inhibits p53 function by physical interaction and phosphorylation. **J. Biol. Chem.**, 2004, **279**:25549-25561.
60. Kato C, Miyazaki K, Nakagawa A, Ohira M, Nakamura Y, **Ozaki T**, Imai T, Nakagawara A. High expression of human tubulin tyrosine ligase and enhanced tubulin tyrosination/detyrosination cycle are associated with neuronal differentiation in neuroblastomas with favorable prognosis. **Int. J. Cancer**, 2004, **112**:365-375.
61. Takahashi M, **Ozaki T**, Todo S, Nakagawara A. Decreased expression of the candidate tumor suppressor gene *ING1* is associated with poor prognosis in advanced neuroblastomas. **Oncology Rep.**, 2004, **12**:811-816.
62. Kramer S, **Ozaki T**, Miyazaki K, Kato C, Hanamoto T, Nakagawara A. Protein stability and function of p73 are modulated by a physical interaction with RanBPM in mammalian cultured cells. **Oncogene**, 2005, **24**:938-944.
63. Lin L, **Ozaki T**, Takada Y, Kageyama H, Nakamura Y, Hata A, Zhang J-H, Simonds W, Nakagawara A, Koseki H. Topors, a p53 and topoisomerase I-binding RING finger protein, is a co-activator of p53 in growth suppression induced by DNA damage. **Oncogene**, 2005, **24**:3385-3396.
64. Hanamoto T, **Ozaki T**, Furuya K, Hosoda M, Hayashi S, Nakanishi M, Yamamoto H, Kikuchi H, Todo S, Nakagawara A. Identification of protein kinase A catalytic subunit b (PKA-Cb) as a novel binding partner of p73 and regulation of p73 function. **J. Biol. Chem.**, 2005, **280**:16665-16675.
65. Aoyama M, **Ozaki T**, Inuzuka H, Tomotsune D, Hirato J, Okamoto Y, Tokita H, Ohira M, Nakagawara A. LMO3 interacts with neuronal transcription factor, HEN2, and acts as an oncogene in neuroblastoma. **Cancer Res.** 2005; **65**: 4587-4597.
66. Hosoda M, **Ozaki T**, Miyazaki K, Hayashi S, Furuya K, Watanabe K, Nakagawa T, Hanamoto T, Todo S, Nakagawara A. UFD2a mediates the proteasomal turnover of p73 without promoting p73 ubiquitination. **Oncogene**, 2005, **24**:7156-7169.
67. Niizuma H, Nakamura Y, **Ozaki T**, Ohira M, Isogai E, Kageyama H, Imaizumi M, Nakagawara A. Bcl-2 is a key regulator for the retinoic acid-induced apoptotic cell death in neuroblastoma. **Oncogene**, 2006, **25**: 5046-5055.
68. Kikuchi H, **Ozaki T**, Furuya K, Hanamoto T, Nakanishi M, Yamamoto H, Yoshida K, Todo S, Nakagawara A. NF- κ B regulates the stability and activity of p73 by inducing its proteolytic degradation through a ubiquitin-dependent proteasome pathway. **Oncogene**, 2006, **25**: 7608-7617.

69. Hayashi S, Ozaki T, Yoshida K, Hosoda M, Todo S, Akiyama S, Nakagawara A. p73 and MDM2 confer the resistance of epidermoid carcinoma to cisplatin by blocking p53. **Biochem. Biophys. Res. Commun.**, 2006, **347**: 60-66.
70. Sang M, Li Y, Ozaki T, Ono S, Ando K, Yamamoto H, Koda T, Geng C, Nakagawara A. p73-dependent induction of 14-3-3s increases the chemo-sensitivity of drug-resistant human breast cancers. **Biochem. Biophys. Res. Commun.**, 2006, **347**: 327-333.
71. Ozaki T, Li Y, Miyazaki K, Tomita T, Iwatsubo T, Nakagawara A. The intracellular domain of the amyloid precursor protein (AICD) enhances the p53-mediated apoptosis. **Biochem. Biophys. Res. Commun.**, 2006, **351**: 57-63.
72. Yamamoto H, Ozaki T, Nakanishi M, Kikuchi H, Yoshida K, Horie H, Kuwano H, Nakagawara A. Oxidative stress induces p53-dependent apoptosis in hepatoblastoma cell through its nuclear translocation. **Genes Cells**, 2007, **12**: 461-471.
73. Nakamura Y, Ozaki T, Niizuma H, Ohira M, Kamijo T, Nakagawara A. Functional characterization of a new p53 mutant generated by homozygous deletion in a neuroblastoma cell line. **Biochem. Biophys. Res. Commun.**, 2007, **354**: 892-898.
74. Takahashi M, Ozaki T, Takahashi A, Miyauchi M, Takada N, Ono S, Koda T, Todo S, Kamijo T, Nakagawara A. DFF45/ICAD restores cisplatin-induced nuclear fragmentation but not DNA cleavage in DFF45-deficient neuroblastoma cells. **Oncogene**, 2007, **26**: 5669-5673.
75. Nakanishi H, Ozaki T, Nakamura Y, Hashizume K, Iwanaka T, Nakagawara A. Purification of human primary neuroblastomas by magnetic beads and their in vitro culture. **Oncology Rep.**, 2007, **17**: 1315-1320.
76. Furuya K, Ozaki T, Hanamoto T, Hosoda M, Hayashi S, Barker PA, Takano K, Matsumoto M, Nakagawara A. Stabilization of p73 by nuclear IKK- α mediates cisplatin-induced apoptosis. **J. Biol. Chem.**, 2007, **282**: 18365-18378.
77. Nakanishi M, Ozaki T, Yamamoto H, Hanamoto T, Kikuchi H, Furuya K, Asaka M, Delia D, Nakagawara A. NFBD1/MDC1 associates with p53 and regulates its function at the crossroad between cell survival and death in response to DNA damage. **J. Biol. Chem.**, 2007, **282**: 22293-23004.
78. Antonelli A, Lenzi L, Nakagawara A, Ozaki T, Chiaretti A, Aloe L. Tumor suppressor proteins are differentially affected in human ependymoblastoma and medulloblastoma cells exposed to nerve growth factor. **Cancer Invest.**, 2007, **25**: 94-101.
79. Ichikawa T, Suenaga Y, Koda T, Ozaki T, Nakagawara A. TAp63-dependent induction of growth differentiation factor 15 (GDF15) plays a critical role in the regulation of keratinocyte differentiation. **Oncogene**, 2008, **27**: 409-420.
80. Yoshida K, Ozaki T, Furuya K, Nakanishi M, Yamamoto H, Ono S, Koda T, Omura K, Nakagawara A. ATM-dependent nuclear accumulation of IKK- α plays an important role in the regulation of p73-mediated apoptosis in response to cisplatin. **Oncogene**, 2008, **27**: 1183-1188.
81. Bu Y, Suenaga Y, Ono, Koda T, Song F, Nakagawara A,

- Ozaki T.** Spl-mediated transcriptional regulation of *NFBD1/MDC1* plays a critical role in DNA damage response pathway **Genes Cells**, 2008, **13**: 55-66.
82. Arai H, **Ozaki T.**, Niizuma H, Nakamura Y, Ohira M, Takano K, Matsumoto M, Nakagawara A. ERAP140/Nbla10993 is a novel favorable prognostic indicator for neuroblastoma and induced in response to retinoic acid. **Oncology Rep.**, 2008, **19**: 1381-1388
 83. Okoshi R, **Ozaki T.**, Yamamoto H, Ando K, Koida N, Ono S, Koda T, Kamijo T, Nakagawara A, Kizaki K. Activation of AMP-activated protein kinase (AMPK) induces p53-dependent apoptotic cell death in response to energetic stress. **J. Biol. Chem.**, 2008, **283**:3979-3987.
 84. Li Y, **Ozaki T.**, Kikuchi H, Yamamoto H, Ohira M, Nakagawara A. A novel HECT-type E3 ubiquitin protein ligase NEDL1 enhances the p53-mediated apoptotic cell death in its catalytic activity-independent manner. **Oncogene**, 2008, **27**: 3700-3709.
 85. Koida N, **Ozaki T.**, Yamamoto H, Ono S, Koda T, Ando K, Okoshi R, Kamijo T, Omura K, Nakagawara A. Inhibitory role of Plk1 in the regulation of p73-dependent apoptosis through physical interaction and phosphorylation. **J. Biol. Chem.**, 2008, **283**: 8555-8563.
 86. Ichikawa T, Suenaga Y, Koda T, **Ozaki T.**, Nakagawara A. DNp63/BMP-7-dependent expression of matrilin-2 is involved in keratinocyte migration in response to wounding. **Biochem. Biophys. Res. Commun.**, 2008, **369**: 994-1000.
 87. Wang H, **Ozaki T.**, Hossain MS, Nakamura Y, Kamij T, Xue X Nakagawara A. A newly identified dependence receptor UNC5H4 is induced during DNA damage-mediated apoptosis and transcriptional target of tumor suppressor p53. **Biochem. Biophys. Res. Commun.**, 2008, **370**: 594-598.
 88. Inoue K, Nakanishi M, Kikuchi H, Yamamoto H, Todo S, Nakagawara A, **Ozaki T.** NFBD1/MDC1 stabilizes oncogenic MDM2 to contribute to cell fate determination in response to DNA damage. **Biochem. Biophys. Res. Commun.**, 2008, **371**: 829-833.
 89. Ando K, Ohira M, **Ozaki T.**, Nakagawa A, Akazawa K, Suenaga Y, Nakamura Y, Koda T, Kamijo T, Murakami Y, Nakagawara A. Expression of *TSLC1*, a candidate tumor suppressor gene mapped to chromosome 11q23, is down-regulated in unfavorable neuroblastoma without promoter hypermethylation. **Int. J. Cancer**, 2008, **123**: 2087-2094.
 90. Hossain MS, **Ozaki T.**, Wang H, Nakagawa A, Takenobu H, Ohira M, Kamijo T, Nakagawara A. N-MYC promotes cell proliferation through a direct transactivation of *neuronal leucine-rich repeat protein-1 (NLRRI)* gene in neuroblastoma. **Oncogene**, 2008, **27**: 6075-6082.
 91. Munirajan AK, Ando K, Mukai A, Takahashi M, Suenaga Y, Ohira M, Koda T, Hirota T, **Ozaki T.**, Nakagawara A. KIF1Bb functions as a haploidinsufficient tumor suppressor gene mapped to chromosome 1p36.2 by inducing apoptotic cell death. **J. Biol. Chem.**, 2008, **283**: 24426-24434.
 92. Yu M, Ohira M, Li Y, Niizuma H, Oo ML, Zhu Y, **Ozaki T.**, Isogai E, Nakamura Y, Koda T, Oba S, Yu B, Nakagawara A.

High expression of *ncRAN*, a novel non-coding RNA mapped to chromosoma 17q25.1, is associated with poor prognosis in neuroblastoma. **Int. J. Oncol.**, in press.

総説その他

1. 岩渕雅樹、滝谷重治、高橋和展、尾崎俊文 (1985). 無細胞転写系における細胞性粘菌アクチン遺伝子のプロモーター解析. **蛋白質核酸酵素** **30**: 126-143.
2. 尾崎俊文、榎本秀樹 (1995). 新規がん抑制遺伝子の構造及び機能解. **Bio Clinica**, **10**: 52-55.
3. 崎山樹、尾崎俊文 (1997). *DAN* 遺伝子. **癌と化学療法** **24**, 1465-1470.
4. 尾崎俊文、中川原章 (2000). p53 ファミリー (わかる細胞周期と癌). **実験医学** 52-60.
5. 尾崎俊文、中川原章 (2000). p73 と p53. **血液腫瘍科** **41**: 332-339.
6. **Ozaki, T.** and Nakagawara, A. (2001). p73 and p53: Functional implication of their interaction. **Recent Res. Dev. Cancer**, **3**: 89-110.
7. 尾崎俊文、中川原章 (2001). BRCT ドメイン (モチーフドメインリスト) 生体の科学 **52**: 519-520.
8. 安藤清宏、尾崎俊文、福澤正洋、中川原章 (2003) p53 の転写活性化能に対する細胞周期制御因子 P1k1 の抑制効果. **日大医学雑誌** **62**: 496-501.
9. 尾崎俊文、中川原章 (2003). p53 ファミリー p63, p73. **Surgery Frontier**, **10**, 64-68.
10. **Ozaki, T.**, Hosoda, M., Miyazaki, K., Hayashi, Y., Watanabe, K., Nakagawa, T. and Nakagawara, A. Functional implication of p73 protein stability in neuronal cell survival and death. (2005). **Cancer Lett.**, **228**: 29-35.
11. **Ozaki, T.** and Nakagawara, A. p73, a sophisticated p53 family member in the cancer world. (2005) **Cancer Sci.**, **96**: 729-737.
12. 古屋武史、上條岳彦、尾崎俊文、草深竹志、中川原章 (2006) 神経芽腫における Shf の機能的役割に関する研究. **日大医学雑誌** **65**: 367-375.
13. 山本英輝、安藤清宏、尾崎俊文、中川原章、桑野博行 (2007) 肝芽腫で高発現するセリンスレオニンキナーゼ P1k1 の機能解析. **The KITAKANTO Medical Journal**, **57**: 82.
14. 尾崎俊文、中川原章 (2008). ATM/IKK- α /p73 経路: 新たな p53 非依存性アポトーシス誘導制御. **生化学** **80**: 409-413.