

**Original Publications** (Miki Ohira, Ph.D.) 1991-2008, 86報

2009 年

Yu M, **Ohira M**, Li Y, Niizuma H, Oo ML, Zhu Y, Ozaki T, Isogai E, Kamijo T, Nakamura Y, Koda T, Oba S, Yu B, Nakagawara A. High expression of *ncRAN*, a novel non-coding RNA mapped to 17q25.1, is associated with poor prognosis in neuroblastoma. *Int. J. Oncol.* *In press*, 2008.

2008 年

Honda S, Arai Y, Haruta M, Sasaki F, **Ohira M**, Yamaoka H, Horie H, Nakagawara A, Hiyama E, Todo S, Kaneko Y. Loss of imprinting of *IGF2* correlates with hypermethylation of the *H19* differentially methylated region in hepatoblastoma. *Br. J. Cancer* 99(11): 1891-1899, 2008.

Ikematsu S, Nakagawara A, Nakamura Y, **Ohira M**, Shinjo M, Kishida S, Kadomatsu K. Plasma midkine level is a prognostic factor for human neuroblastoma. *Cancer Sci.* 99(10): 2070-2074, 2008.

Miyake I, **Ohira M**, Nakagawara A, Sakai R. Distinct role of ShcC docking protein in the differentiation of neuroblastoma. *Oncogene* 28(5): 662-673, 2009.

Chen Y, Takita J, Choi YL, Kato M, **Ohira M**, Sanada M, Wang L, Soda M, Kikuchi A, Igarashi T, Nakagawara A, Hayashi Y, Mano H, Ogawa A. Novel oncogenic mutations of ALK kinase in neuroblastoma. *Nature* 455(7215): 971-4, 2008.

Munirajan AK, Ando K, Mukai A, Takahashi M, Suenaga Y, **Ohira M**, Koda T, Hirota T, Ozaki T, Nakagawara A. KIF1Bbeta functions as a haploidinsufficient tumor suppressor gene mapped to chromosome 1p36.2 by inducing apoptotic cell death. *J Biol Chem.* 283(36): 24426-34, 2008.

Hossain S, Ozaki T, Wang H, Nakagawa A, Takenobu H, **Ohira M**, Kamijo T, Nakagawara A. N-MYC promotes cell proliferation through a direct transactivation of neuronal leucine-rich repeat protein-1 (NLRR1) gene in neuroblastoma. *Oncogene* 27(46): 6075-6082, 2008.

Ando K, **Ohira M**, Ozaki T, Nakagawa A, Akazawa K, Suenaga Y, Nakamura Y, Koda T, Kamijo T, Murakami Y, Nakagawara A. Expression of *TSLC1*, a candidate tumor suppressor gene mapped to chromosome 11q23, is down-regulated in unfavorable neuroblastoma without promoter hypermethylation. *Int. J. Cancer* 123(9): 2087-94, 2008.

Nakagawa H, **Ohira M**, Hayashi S, Abe S, Saito S, Nagahori N, Monde K, Shinohara Y, Fujitani N, Kondo H, Akiyama S, Nakagawara A, Nishimura S. Alterations in the glycoform of cisplatin-resistant human carcinoma cells are caused by defects in the endoplasmic reticulum-associated degradation system. *Cancer Lett* 270(2): 295-301, 2008.

Honda S, Haruta M, Sugawara W, Sasaki F, **Ohira M**, Matsunaga T, Yamaoka H, Horie H, Ohnuma N, Nakagawara A, Hiyama E, Todo S, Kaneko Y. The methylation

status of RASSF1 promoter predicts responsiveness to chemotherapy and eventual cure in hepatoblastoma patients. *Int J Cancer* 123(5): 1117-25, 2008.

Abe M, Watanabe N, McDonnell N, Takato T, **Ohira M**, Nakagawara A, Ushijima T. Identification of genes targeted by CpG island methylator phenotype in neuroblastomas, and their possible integrative involvement in poor prognosis. *Oncology* 74(1-2): 50-60, 2008.

Li Y, Ozaki T, Kikuchi H, Yamamoto H, **Ohira M**, Nakagawara A. A novel HECT-type E3 ubiquitin protein ligase NEDL1 enhances the p53-mediated apoptotic cell death in its catalytic activity-independent manner. *Oncogene* 27, 3700-9, 2008.

Arai H, Ozaki T, Niizuma H, Nakamura Y, **Ohira M**, Takano K, Matsumoto M, Nakagawara A. ERAP140/Nbla10993 is a novel favorable prognostic indicator for neuroblastoma induced in response to retinoic acid. *Oncology Rep.* 19(6): 1381-8, 2008. (Jun)

Kurata K, Yanagisawa R, **Ohira M**, Kitagawa M, Nakagawara A, Kamijo T. Stress via p53 pathway causes apoptosis by mitochondrial Noxa up-regulation in doxorubicin-treated neuroblastoma cells. *Oncogene* 27(6): 741-754, 2008.

Tomioka N, Oba S, **Ohira M**, Misra A, Fridlyand J, Ishii S, Nakamura Y, Isogai E, Hirata T, Yoshida Y, Todo S, Kaneko Y, Albertson D, Pinkel D, Feuerstein B, Nakagawara A. Novel risk stratification of patients with neuroblastoma by genomic signature which is independent of molecular signature. *Oncogene* 27, 441-449, 2008.

2007 年

Nakamura Y, Ozaki T, Niizuma H, **Ohira M**, Kamijo T, Nakagawara A. Functional characterization of a new p53 mutant generated by homozygous deletion in a neuroblastoma cell line. *Biochem. Biophys. Res. Commun.* 354, 892-898, 2007. Mar 23;354(4):892-8.

Kaneko S, **Ohira M**, Nakamura Y, Isogai E, Nakagawara A, Kaneko M. Relationship of DDX1 and NAG gene amplification/overexpression to the prognosis of patients with MYCN-amplified neuroblastoma. *J Cancer Res Clin Oncol.* 133, 185-192, 2007.

2006 年

Takenaga K, Nygren J, Zelenina M, **Ohira M**, Iuchi T, Lukanidin E, Sjoquist M, Kozlova NE. Modified expression of S100A4/Mts1 protein in C6 glioma cells or surrounding astrocytes affects migration of tumor cells *in vitro* and *in vivo*. *Neurobiol Dis* 25, 455-463, 2007.

Oba S, Tomioka N, **Ohira M**, Ishii S. Combfit: a normalization method for array CGH data. *IPSI Transactions on Bioinformatics* 47, 73-82, 2006.

Niizuma H, Nakamura Y, Ozaki T, **Ohira M**, Isogai E, Kageyama H, Imaizumi M, Nakagawara A. Bcl-2 is a key regulator for the retinoic acid-induced apoptotic cell death in neuroblastoma. *Oncogene* 25(36):5046-55, 2006.

Machida T, Fujita T, Oo M L, **Ohira M**, Isogai E, Mihara M, Hirato J, Tomotsune D, Hirata T, Fujimori M, Adachi W, Nakagawara A. Increased expression of pro-apoptotic *BMCC1*, a novel gene with the *BNIP2* and *Cdc42GAP* Homology (BCH) domain, is associated with favorable prognosis in human neuroblastomas. *Oncogene* 25, 1931-42, 2006.

Inamori K, Gu J, **Ohira M**, Kawasaki A, Nakamura Y, Nakagawa T, Kondo A, Miyoshi E, Nakagawara A, Taniguchi N. High expression of N-acetylglucosaminyltransferase V in favorable neuroblastomas: Involvement of its effect on apoptosis. *FEBS Lett.* 580, 627-32, 2006.

Chen H, Suzuki M, Nakamura Y, **Ohira M**, Ando S, Iida T., Nakajima T, Nakagawara A, Kimura H. Aberrant methylation of *RASGRF2* and *RASSF1A* in human non-small cell lung cancer. *Oncol. Rep.* 15, 1281-5, 2006.

2005 年

Osajima-Hakomori Y, Miyake I, **Ohira M**, Nakagawara A, Nakagawa A, Sakai R. Biological role of anaplastic lymphoma kinase in neuroblastoma. *Am. J. Pathol.* 167, 213-22, 2005.

Chen H, Suzuki M, Nakamura Y, **Ohira M**, Suenaga Y, Ando S, Iida T., Nakajima T, Nakagawara A, Kimura H. Aberrant methylation of *FBN2* correlates with progression of human non-small cell lung cancer. *Lung Cancer* 50, 43-9, 2005.

Gotoh T, Hosoi H, Iehara T, Kuwahara Y, Osonoe S, Tsuchiya K, Kuroda H, **Ohira M**, Nakagawara A, Sugimoto T. Prediction of *MYCN* amplification in neuroblastoma using serum DNA and real-time quantitative PCR. *J. Clin. Oncol.* 23, 5205-10, 2005.

**Ohira M**, Oba S, Nakamura Y, Isogai E, Kaneko S, Nakagawa A, Hirata T, Kubo H, Goto T, Yamada S, Yoshida Y, Fuchioka M, Ishii S, Nakagawara A. Expression profiling using a tumor-specific cDNA microarray predicts the prognosis of intermediate-risk neuroblastomas. *Cancer Cell*, 7, 337-50, 2005.

Aoyama M, Ozaki T, Inuzuka H, Tomotsune D, Hirato J, Okamoto Y, Tokita H, **Ohira M**, Nakagawara A. LMO3 interacts with neuronal transcription factor, *HEN2*, and acts as an oncogene in neuroblastoma. *Cancer Res.* 65, 4587-97, 2005.

**Ohira M**, Oba S, Nakamura Y, Hirata T, Ishii S, Nakagawara A. A review of DNA microarray analysis of human neuroblastomas. *Cancer Lett.*, 228, 5-11, 2005.

Abe M, **Ohira M**, Kaneda A, Yagi Y, Yamamoto S, Kitano Y, Takato T, Nakagawara A, Ushijima T. CpG island methylator phenotype is a strong determinant of poor prognosis in neuroblastomas. *Cancer Res.* 65, 828-34, 2005.

2004 年

Kato C, Miyazaki K, Nakagawa A, **Ohira M**, Nakamura Y, Ozaki T, Imai T, Nakagawara A. Low expression of human tubulin tyrosine ligase and suppressed tubulin tyrosination/detyrosination cycle are associated with impaired neuronal differentiation in neuroblastomas with poor prognosis. *Int. J. Cancer* 112, 365-75, 2004.

Yamada S, **Ohira M**, Horie H, Ando K, Takayasu H, Suzuki Y, Sugano S, Hirata T, Goto T, Matsunaga T, Hiyama E, Hayashi Y, Ando H, Suita S, Kaneko M, Sasaki F, Hashizume K, Ohnuma N, Nakagawara A. Expression profiling and differential screening between hepatoblastomas and the corresponding normal livers: Identification of high expression of the *PLK1* oncogene as a poor-prognostic indicator of hepatoblastomas. *Oncogene* 23, 5901-11, 2004.

Hamano S, **Ohira M**, Isogai E, Nakada K, Nakagawara A. Identification of novel human neuronal leucine-rich repeat (hNLR) family genes and inverse association of expression of *Nbla10449/hNLR-1* and *Nbla10677/hNLR-3* with the prognosis of primary neuroblastomas. *Int. J. Oncology*, 24, 1457-66, 2004.

Nakagawara A, **Ohira M**. Comprehensive genomics linking between neural development and cancer: neuroblastoma as a model. *Cancer Lett.* 204, 213-24, 2004.

2003 年

**Ohira M**, Morohashi A, Inuzuka H, Shishikura T, Kawamoto T, Kageyama H, Nakamura Y, Isogai E, Takayasu H, Sakiyama S, Suzuki Y, Sugano S, Goto T, Sato S, Nakagawara A. Expression profiling and characterization of 4200 genes cloned from primary neuroblastomas: identification of 305 genes differentially expressed between favorable and unfavorable subsets. *Oncogene* 22, 5525-36, 2003.

**Ohira M**, Morohashi A, Nakamura Y, Isogai E, Furuya K, Hamano S, Machida T, Aoyama M, Fukumura M, Miyazaki K, Suzuki Y, Sugano S, Hirato J, Nakagawara A. Neuroblastoma oligo-capping cDNA project: toward the understanding of the genesis and biology of neuroblastoma. *Cancer Lett.*, 197, 63-8, 2003.

Kawamoto T, **Ohira M**, Hamano S, Hori T, Nakagawara A. High expression of the novel endothelin-converting enzyme genes, *Nbla03145/ECEL1 $\alpha$*  and  $\beta$ , is associated with favorable prognosis in human neuroblastomas. *Int. J. Oncology*, 22, 815-22, 2003.

Tomioka N, Kobayashi H, Kageyama H, **Ohira M**, Nakamura Y, Sasaki F, Todo S, Nakagawara A, Kaneko Y. Chromosomes that show partial loss or gain in near-diploid tumors coincide with chromosomes that show whole loss or gain in near-triploid tumors: Evidence suggesting the involvement of the same genes in the tumorigenesis of high- and low-risk neuroblastomas. *Genes Chromosomes Cancer*. 36, 139-50, 2003.

2002 年

Takahashi M, Seki N, Ozaki T, Kato M, Kuno T, Nakagawa T, Watanabe K, Miyazaki K, **Ohira M**, Hayashi S, Hosoda M, Tokita H, Mizuguchi H, Hayakawa T, Todo S, Nakagawara A. Identification of the p33ING1-regulated genes which include cyclin B1 and proto-oncogene DEK by using cDNA microarray in a mouse mammary epithelial cell line NMuMG. *Cancer Res.* 62, 2203-9, 2002.

Iuchi T, Namba H, Iwadate Y, Shishikura T, Kageyama H, Nakamura Y, **Ohira M**, Tamaura A, Osato K, Sakiyama S, Nakagawara A. Identification of the small interstitial deletion at chromosome 1p34-p35 and its association with poor outcome in oligodendroglial tumors. *Genes Chromosomes Cancer*. 35, 170-5, 2002.

2001 年

Yang X, Aoki Y, Li X, Sakamoto O, Hiratsuka M, Kure S, Taheri S, Christensen E, Inui K, Kubota M, **Ohira M**, Ohki M, Kudoh J, Kawasaki K, Shibuya K, Shintani A, Asakawa S, Minoshima S, Shimizu N, Narisawa K, Matsubara Y, Suzuki Y. Structure of human holocarboxylase synthetase gene and mutation spectrum of holocarboxylase synthetase deficiency. *Hum Genet.* 109, 526-534, 2001.

Shinagawa T, Nomura T, Colmenares C, **Ohira M**, Nakagawara A, Ishii S. Increased susceptibility to tumorigenesis of ski-deficient heterozygous mice. *Oncogene* 20, 8100-8, 2001.

Ichimiya S, Nimura Y, Kageyama H, Takada N, Sunahara M, Shishikura T, Nakamura Y, Sakiyama S, Seki N, **Ohira M**, Kaneko Y, McKeon F, Caput D, Nakagawara A. Genetic analysis of p73 localized at chromosome 1p36.3 in primary neuroblastomas. *Med Pediatr Oncol.* 36, 42-4, 2001.

Chen YZ, Hayashi Y, Wu JG, Takaoka Ei, Maekawa K, Watanabe N, Inazawa J, Hosoda F, Arai Y, Ohki M, Mizushima H, Morohashi A, **Ohira M**, Nakagawara A, Liu SY, Hoshi M, Horii A, Soeda E. A bac-based sts-content map spanning a 35-mb region of human chromosome 1p35-p36. *Genomics* 74, 55-70, 2001.

2000 年

**Ohira M**, Shishikura T, Kawamoto T, Inuzuka H, Morohashi A, Takayasu H, Kageyama H, Takada N, Takahashi M, Sakiyama S, Suzuki Y, Sugano S, Kuma H, Nozawa I, Nakagawara A. Hunting the subset-specific genes of neuroblastoma: expression profiling and differential screening of the full-length-enriched oligo-capping cDNA libraries. *Med. Pediatr. Oncol.* 35, 547-9, 2000.

Nakagawara A, **Ohira M**, Kageyama H, Mihara M, Furuta S, Machida T, Takayasu H, Islam A, Nakamura Y, Takahashi M, Shishikura T, Kaneko Y, Toyoda A, Hattori M, Sakaki Y, Ohki M, Horii A, Soeda E, Inazawa J, Seki N, Kuma H, Nozawa I, Sakiyama S. Identification of the homozygously deleted region at chromosome 1p36.2 in human neuroblastoma. *Med. Pediatr. Oncol.* 35, 516-21, 2000.

Furuta S, **Ohira M**, Machida T, Hamano S, Nakagawara A. Analysis of loss of heterozygosity at 16p12-p13 (familial neuroblastoma locus) in 470 neuroblastomas including both sporadic and mass screening tumors. *Med. Pediatr. Oncol.* 35, 531-3, 2000.

Kawamoto T, Shishikura T, **Ohira M**, Takayasu H, Morohashi A, Takada N, Takahashi M, Suzuki Y, Sugano S, Hori T, Nakagawara A. Association between favorable neuroblastoma and high expression of the novel metalloproteinase gene, *nbla3145/XCE*, cloned by differential screening of the full-length-enriched oligo-capping neuroblastoma cDNA libraries. *Med. Pediatr. Oncol.* 35, 628-31, 2000.

**Ohira M**, Kageyama H, Mihara M, Furuta S, Machida T, Shishikura T, Takayasu H, Islam A, Nakamura Y, Takahashi M, Tomioka N, Sakiyama S, Kaneko Y, Toyoda A, Hattori M, Sakaki Y, Ohki M, Horii A, Soeda E, Inazawa J, Seki N, Kuma H, Nozawa I, Nakagawara A. Identification and characterization of a 500-kb homozygously deleted region at 1p36.2-p36.3 in a neuroblastoma cell line. *Oncogene* 19, 4302-7, 2000.

Hanaoka E, Ozaki T, **Ohira M**, Nakamura Y, Suzuki M, Takahashi E, Moriya H, Nakagawara A, Sakiyama S. Molecular cloning and expression analysis of the

human DA41 gene and its mapping to chromosome 9q21.2-q21.3. *J Hum Genet.* 45, 188-91, 2000.

Nagai M, Ichimiya S, Ozaki T, Seki N, Mihara M, Furuta S, **Ohira M**, Tomioka N, Nomura N, Sakiyama S, Kubo O, Takakura K, Hori T, and Nakagawara A. Identification of the full-length KIAA0591 gene encoding a novel kinesin-related protein which is mapped to the neuroblastoma suppressor gene locus at 1p36.2. *Int. J. Oncol.* 16, 907-16, 2000.

Islam A, Kageyama H, Takada N, Kawamoto T, Takayasu H, Isogai E, **Ohira M**, Hashizume K, Kobayashi H, Kaneko Y, and Nakagawara A. High expression of Survivin, mapped to 17q25, is significantly associated with poor prognostic factors and promotes cell survival in human neuroblastoma. *Oncogene* 19, 617-23, 2000.

Ozaki, T., Nagase, T., Ichimiya, S., Seki, N., **Ohira, M.**, Nomura, N., Takada, N., Sakiyama, S., Weber, B.L., and Nakagawara, A. NFBD1/KIAA0170 is a novel nuclear transcriptional transactivator with BRCT domain. *DNA Cell Biol.* 19, 475-85, 2000.

1999 年

Seki, N., Hayashi, A., Hattori, A., Kozuma, S., **Ohira, M.**, Hori, T., and Saito, T. Chromosomal assignment of a human apoptosis-associated tyrosine kinase gene on chromosome 17q25.3 by somatic hybrid analysis and fluorescence in situ hybridization. *J. Hum. Genet.* 44, 141-142, 1999.

Ueki, N., Seki, N., Yano, K., **Ohira, M.**, Saito, T., Masuho, Y., and Muramatsu, M. Isolation and chromosomal assignment of human genes encoding cofactor of LIM homeodomain proteins, CLIM1 and CLIM2. *J. Hum. Genet.* 44, 112-115, 1999.

Ichimiya, S., Nimura, Y., Kageyama, H., Takada, N., Sunahara, M., Shishikura, T., Nakamura, Y., Sakiyama, S., Seki, N., **Ohira, M.**, Kaneko, Y., McKeon, F., Caput, D., and Nakagawara, A. p73 at chromosome 1p36.3 is lost in advanced stage neuroblastoma but its mutation is infrequent. *Oncogene* 18, 1061-1066, 1999.

Seki, N., Hattori, A., Hayashi, A., Kozuma, S., **Ohira, M.**, Hori, T., Saito, T. Structure, expression profile and chromosomal location of an isolog of DNA-PKcs interacting protein (KIP) gene1. *Biochim. Biophys. Acta* 1444, 143-147, 1999.

1998 年

Nimura, Y., Mihara, M., Ichimiya, S., Sakiyama, S., Seki, N., **Ohira, M.**, Nomura, N., Fujimori, M., Adachi, W., Amano, J., He, M., Ping, Y.M., Nakagawara, A. p73, a gene related to p53, is not mutated in esophageal carcinomas. *Int. J. Cancer*, 78, 437-440, 1998.

Seki, N., Hayashi, A., Abe, M., Araki, R., Fujimori, A., Fukumura, R., Hattori, A., Kozuma, S., **Ohira, M.**, Hori, T., Saito, T. Chromosomal assignment of the gene for human DNA-PKcs interacting protein (KIP) on chromosome 15q25.3-q26.1 by somatic hybrid analysis and fluorescence in situ hybridization. *J. Hum. Genet.* 43, 275-277, 1998.

Seki, N., Muramatsu, M., Sugano, S., Suzuki, Y., Nakagawara, A., **Ohira, M.**, Hayashi,

A., Hori, T., Saito, T. Cloning, expression analysis, and chromosomal localization of *HIP1R*, an isolog of huntingtin interacting protein (HIP1). *J. Hum. Genet.* 43, 268-271, 1998.

Seki, N., Hattori, A., Sugano, S., Suzuki, Y., Nakagawara, A., **Ohira, M.**, Muramatsu, M., Hori, T., Saito, T. Isolation, tissue expression, and chromosomal assignment of a novel human gene which encodes a protein with RING finger motif. *J. Hum. Genet.* 43, 272-274, 1998.

Saito, T., Seki, N., **Ohira, M.**, Hayashi, A., Kozuma, S., Hattori, A., Hori, T. Assignment of the ZIP kinase gene to human chromosome 19p13.3 by somatic hybrid analysis and fluorescence in-situ hybridization. *J. Hum. Genet.* 43, 209-211, 1998.

Seki, N., Sugano, S., Suzuki, Y., Nakagawara, A., **Ohira, M.**, Muramatsu, M., Saito, T., Hori, T. Isolation, tissue expression, and chromosomal assignment of human RGS5, a novel G-protein signaling regulator gene. *J. Hum. Genet.* 43, 202-205, 1998.

Senoo, M., Seki, N., **Ohira, M.**, Sugano, S., Watanabe, M., Tachibana, M., Tanaka, T., Shinkai, Y., Kato, H. A second p53-related protein, p73L, with high homology to p73. *Biochem. Biophys. Res. Commun.* 248, 603-607, 1998.

**Ohira, M.**, Seki, N., Nagase, T., Ishikawa, K., Nomura, N. and Ohara, O. Characterization of a human homolog of the mouse *Bach1* gene encoding a BTB-basic leucine zipper transcription factor and its mapping to chromosome 21q22.1. *Genomics* 47, 300-306, 1998.

Hori, T., Seki, N., **Ohira, M.**, Saito, T., Yamauchi, M., Sagara, M., Hayashi, A., Tsuji, S., Ito, H. and Imai, T. A distamycin-inducible fragile site, FRA8E, is located in the region of the hereditary multiple exostoses gene and is not involved in HPV16 DNA integration and amplification. *Cancer genet. Cytogenet.* 101, 24-34, 1998.

1997 年

Korenberg, J. R., Aaltonen, J., Brahe, C., Cabin, D., Creau, N., Delabar, J. M., Doering, J., Gardiner, K., Hubert, R. S., Ives, J., Kessling, A., Kudoh, J., Lafrenière, Murakami, Y., **Ohira, M.**, Ohki, M., Patterson, D., Potier, M. C., Quackenbush, J., Reeves, R. H., Sakaki, Y., Shimizu, N., Soeda, E., Van Broeckhoven, C., and Yaspo, M. L. Report of the Sixth International Workshop on Human Chromosome 21 Mapping 1996. *Cytogenet. Cell Genet.* 79, 21-52, 1997.

Seki, N., **Ohira, M.**, Nagase, T., Ishikawa, K., Miyajima, N., Nakajima, D., Nomura, N., and Ohara, O. Characterization of cDNA clones in size-fractionated cDNA libraries from human brain. *DNA Res.* 4, 345-349, 1997.

Ishikawa, K., Nagase, T., Nakajima, D., Seki, N., **Ohira, M.**, Miyajima, N., Tanaka, A., Kotani, H., Nomura, N. and Ohara, O. Prediction of the coding sequences of unidentified human genes. VIII. 78 new cDNA clones from brain which code for large proteins *in vitro*. *DNA Res.* 4, 307-313, 1997.

Seki, N., Nimura, Y., **Ohira, M.**, Saito, T., Ichimiya, S., Nomura, N. and Nakagawara, A. Identification and chromosome assignment of a human gene encoding a novel phosphatidylinositol-3 kinase. *DNA Res.* 4, 355-358, 1997.

Saito, T., Seki, N., Ishii, H., **Ohira, M.**, Hayashi, A., Kozuma, S. and Hori, T. Complementary DNA cloning and chromosomal mapping of a novel phosphatidylinositol kinase gene. *DNA Res.* 4, 301-305, 1997.

Ozaki, T., Enomoto, H., Nakamura, Y., Kondo, K., Seki, N., **Ohira, M.**, Nomura, N., Ohki, M., Nakagawara, A. and Sakiyama, S. The genomic analysis of human DNA gene. *DNA and Cell Biol.* 16, 1031-1039, 1997.

**Ohira, M.**, Seki, N., Nagase, T., Suzuki, E., Nomura, N., Ohara, O., Hattori, M., Sakaki, Y., Eki, T., Murakami, Y., Saito, T., Ichikawa, H. and Ohki, M. Gene identification in the 1.6-Mb region of the Down syndrome region on chromosome 21. *Genome Res.* 7, 47-58, 1997.

Nagase, T., Ishikawa, K., Nakajima, D., **Ohira, M.**, Seki, N., Miyajima, N., Kotani, H., Nomura, N., and Ohara, O. Prediction of the coding sequences of unidentified human genes. VII. The complete sequences of 100 new cDNA clones from brain which can code for large proteins *in vitro*. *DNA Res.* 4, 141-150, 1997.

Ohara, O., Nagase, T., Ishikawa, K., Nakajima, D., **Ohira, M.**, Seki, N., and Nomura, N. Construction and characterization of human brain cDNA libraries suitable for analysis of cDNA clones encoding relatively large proteins. *DNA Res.* 4, 53-59, 1997.

Saitoh K, Miura I, Ohshima A, Takahashi N, Kume M, Utsumi S, Kobayashi Y, Hashimoto K, Hatano Y, Nimura T, Saito M, Enomoto K, **Ohhira M**, Shimizu K, Ohki M, Miura AB. Translocation (8;12;21)(q22.1;q24.1;q22.1): a new masked type of t(8;21)(q22;q22) in a patient with acute myeloid leukemia. *Cancer Genet Cytogenet.* 96, 111-4, 1997.

1996 年

**Ohira, M.**, Ichikawa, H., Suzuki, E., Iwaki, M., Suzuki, K., Saito-Ohara, F., Ikeuchi, T., Chumakov, I., Tanahashi, H., Tashiro, K., Sakaki, Y., and Ohki, M. A 1.6-Mb P1-based physical map of the Down syndrome region on chromosome 21. *Genomics* 33, 65-74, 1996.

**Ohira, M.**, Ootsuyama, A., Suzuki, E., Ichikawa, H., Seki, N., Nagase, T., Nomura, N., and Ohki, M. Identification of a novel gene containing the tetratricopeptide repeat motifs of the Down syndrome region on chromosome 21. *DNA Res.* 3, 9-16, 1996.

Nagase, T., Seki, N., Ishikawa, K., **Ohira, M.**, Kawarabayashi, Y., Ohara, O., Tanaka, A., Kotani, H., Miyajima, N., and Nomura, N. Prediction of the coding sequences of unidentified human genes. VI. The coding sequences of 80 new genes (KIAA0201-KIAA0280) deduced by analysis of cDNA clones from cell line KG-1 and brain. *DNA Res.* 3, 321-329, 1996.

1995 年

Goto, M., Shimizu, T., Sawada, J., Sawa, C., Watanabe, H., Ichikawa, H., **Ohira, M.**, Ohki, M., and Handa, H. Assignment of the E4TF1-60 gene to human chromosome 21q21.2-q21.3. *Gene* 166, 337-338, 1995.

Miyoshi, H., **Ohira, M.**, Shimizu, K., Mitani, K., Hirai, H., Imai, T., Yokoyama, K., Soeda, E., and Ohki, M. Alternative splicing and genomic structure of the AML1

gene involved in acute myeloid leukemia. *Nucleic Acids Res.* 23, 2762-2769, 1995.

1994 年

Tanahashi, H., Ito, T., Hattori, M., **Ohira, M.**, Ohki, M., Tashiro, K., and Sakaki, Y. Sixty new STSs (sequence-tagged sites) of human chromosome 21. *DNA Res.* 1, 85-89, 1994.

1993 年

Shimizu, K., Ichikawa, H., Tojo, A., Kaneko, Y., Maseki, N., Hayashi, Y., **Ohira, M.**, Asano, S., and Ohki, M. An ets-related gene, ERG, is rearranged in human myeloid leukemia with t(16;21) chromosomal translocation. *Proc. Natl. Acad. Sci. USA* 90, 10280-10284, 1993.

Ichikawa, H., Hosoda, F., Arai, Y., Shimizu, K., **Ohira, M.**, and Ohki, M. A Not I restriction map of the entire long arm of human chromosome 21. *Nat. Genet.* 4, 361-366, 1993.

1991 年

**Ohira, M.**, Bae, Y. S., and Ikeda, H. A new type of insertion mutation in monkey cells: insertion accompanied by long target site duplication. *Mol. Gen. Genet.* 229, 325-333, 1991.

Bae, Y. S., Chiba, M., **Ohira, M.**, and Ikeda, H. A shuttle vector for analysis of illegitimate recombination in mammalian cells: effects of DNA topoisomerase inhibitors on deletion frequency. *Gene* 101, 285-289, 1991.

## 日本語総説 (11 編)

がん診療ハンドブック

監修 竜 崇正 編集 木村秀樹 永井書店

第3部 先進医療

9 がんプロトコールスタディにおける前向き附随研究の推進 p117-119

大平美紀、中川原 章

第4章 DNAチップ／マイクロアレイ臨床応用への実際 2. 個別化医療 5) 神経芽細胞腫

遺伝子医学 MOOK 10号 DNAチップ／マイクロアレイ臨床応用の実際 —基礎・技術、臨床・創薬応用の実際から今後の展開・問題点まで— 編集 油谷浩幸 メディカル ドウ 2008

年 321-327

大平美紀、中川原 章

特集 小児固形腫瘍の分子生物学（その3）：最新の知見

発現アレイと小児固形腫瘍

小児外科 2007 November vol. 39 No. 11 1277-1283

東京医学社

大平美紀、中川原 章

臨床遺伝学'07 ―ゲノム科学の臨床へのインパクト―

ゲノムで進むがん研究 小児がんの網羅的遺伝子発現解析とその臨床応用

最新医学 第62巻／9月増刊号 通巻第780号 2007 9月増刊号 2077-2086

編集 菅野純夫

最新医学社

大平美紀、中川原 章

III 医療での実用化を目指した応用 神経芽腫診断用発現チップの開発と実用化

特集 ゲノム創薬・疾患診断を実現する新世代マイクロアレイ 企画 稲澤譲治

バイオテクノロジージャーナル 2007 5-6 vol. 7 No. 3 308-313

羊土社

大平美紀、中川原 章

新小児がんの診断と治療

別所文雄、杉本徹、横森欣司 編集

分子生物学的概要 ii) マイクロアレイの応用 pp. 38-46, 2007

診断と治療社

大平美紀、中川原 章

特集「固形腫瘍の新しい治療」・小児固形腫瘍病因解明のための基礎的研究

小児科診療 vol. 67, No. 4(13), 529-536, 2004.

中川原 章、大平美紀

特集「ミレニアム医学の構築-固形癌」・神経芽腫の予後予測と遺伝子診断-現在と近未来-

医学のあゆみ vol. 197, No. 13, 1169-1174, 2001.

中川原 章、大平美紀

特集「がんとゲノム 1-発生領域より」・ゲノム研究からみた神経芽細胞腫

ゲノム医学 vol.1, No. 3, 255-264, 2001.

**Ohira, M.**, and Ohki, M Genome analysis of human chromosome 21: transcript mapping of the Down syndrome region. *Seikagaku* **69**, 229-243, 1997.

**Ohira, M.**, Ichikawa, H., and Ohki, M. [Physical mapping of human chromosomes with linking clones and jumping clones] *Nippon Rinsho* **51**, 2252-2257, 1993.